



Pocket Genius

DINOSAURS



FACTS AT YOUR FINGERTIPS



Pocket Genius

DINOSAURS



FACTS AT YOUR FINGERTIPS



Penguin
Random
House

DK DELHI

Project editor Vriken Chopra
Project art editor Nishesh Batnagar
Senior editor Kingshuk Ghoshal
Senior art editor Govind Mittal
Art editor Isha Nagar

DTP designers Mohammad Usman, Vishal Bhatia
Picture researcher Sumedha Chopra

DK LONDON

Senior editor Dr. Rob Houston
Senior art editor Philip Letsu
US editor Margaret Parrish
Jacket editor Manisha Majithia
Jacket designer Laura Brim
Jacket design development manager
Sophia Tampakopolous
Production editor Ben Marcus
Production controller Mary Slater
Publisher Andrew Macintyre
Associate publishing director Liz Wheeler
Art director Phil Ormerod
Publishing director Jonathan Metcalf
Consultant Professor Michael J. Benton

First American Edition, 2012

This edition published in the United States in 2016 by
DK Publishing, 345 Hudson Street, New York, New York 10014

Copyright © 2012, 2016 Dorling Kindersley Limited
DK, a Division of Penguin Random House LLC
16 17 18 19 20 10 9 8 7 6 5 4 3 2 1
001-290704-January/2016

All rights reserved.

Without limiting the rights under the copyright reserved above, no part of this publication may be reproduced, stored in or introduced into a retrieval system, or transmitted, in any form, or by any means (electronic, mechanical, photocopying, recording, or otherwise), without the prior written permission of the copyright owner. Published in Great Britain by Dorling Kindersley Limited.

A catalog record for this book
is available from the Library of Congress.
ISBN: 978-1-4654-4561-2

DK books are available at special discounts when purchased in bulk for sales promotions, premiums, fund-raising, or educational use. For details, contact: DK Publishing Special Markets, 345 Hudson Street, New York, New York 10014
SpecialSales@dk.com

Printed and bound in China

A WORLD OF IDEAS:
SEE ALL THERE IS TO KNOW
www.dk.com



CONTENTS

- 4 Before the dinosaurs
- 6 Dinosaurs and after
- 8 The Triassic
- 10 The Jurassic
- 12 The Cretaceous
- 14 Dinosaur ancestors
- 16 Types of dinosaur
- 18 Dinosaurs to birds
- 20 Dinosaurs die out
- 22 How fossils form
- 24 Fossil bones
- 26 Exceptional fossils
- 28 Trace fossils
- 30 Reconstruction

32 DINOSAURS

- 34 The first dinosaurs
- 36 Theropods
- 54 Early birds
- 56 Prosauropods
- 60 Sauropods
- 72 Stegosaurs and relatives
- 78 Nodosaur
- 80 Ankylosaur
- 84 Ornithomimid
- 96 Pachycephalosaurs
- 98 Ceratopsians

102 DINOSAUR NEIGHBORS

- 104 Rhynchosaurs
- 106 Archosaurs
- 114 Cynodonts and dicynodonts
- 116 Early mammals

118 SEA REPTILES

- 120 Placodonts and turtles
- 122 Nothosaurs
- 124 Ichthyosaurs
- 126 Plesiosaurs
- 128 Pliosaurus
- 132 Mosasaurs

134 FLYING REPTILES

- 136 Pterosaurs
- 142 Prehistoric records
- 144 Largest dinosaurs
- 146 Dinosaur discoveries
- 148 Glossary
- 152 Index
- 156 Acknowledgments

Scales and sizes


This book contains profiles of prehistoric animals with scale drawings to indicate their size.



6 ft
(1.8 m)



6 in
(15 cm)



Kentrosaurus


Before the dinosaurs

The Earth is more than 4.5 billion years old, and the first traces of life appeared some 1 billion years after it formed. Single-celled organisms were the first life-forms to appear, and over millions of years they evolved into invertebrates (animals without backbones) and vertebrates (animals with backbones). The Earth's history is divided into eras, which are further divided into periods.

Cambrian Explosion

Around 530 million years ago (mya), there was a rapid increase in the number of different kinds of invertebrate animals.

This is known as the Cambrian Explosion, because it occurred in the Cambrian Period.



Anomalocaris, one of the top predators of the Cambrian oceans

PRECAMBRIAN

Cambrian
PALEOZOIC ERA

Ordovician

Silurian

542 mya

488 mya


443 mya

Precambrian life

Some of the first living organisms were bacteria living in mats on the seabed.

They trapped sand and used it to build rocky mounds called stromatolites, just as some still do today (left).

Some fossil stromatolites are an incredible 3.5 billion years old.




Stromatolites



Devonian plants

The Devonian Period lasted from 416 to 359 mya. It was in this period that the plant *Archaeopteris* appeared. It was the first tree to form dense forests that spread across the planet.



Dimetrodon

Permian reptiles

Many different reptile groups were widespread in the Permian Period, including ancestors of the dinosaurs. *Dimetrodon*, however, was a remote ancestor of mammals.

Archaeopteris


Devonian

Carboniferous

Permian

299 mya

252 mya




Amphibamus

Carboniferous amphibians

First appearing in the Devonian Period, four-legged vertebrates, or tetrapods, were common on land by the Carboniferous Period. The amphibian *Amphibamus* was an early tetrapod.

Dinosaurs and after

The dinosaurs lived in the Mesozoic Era, which lasted from 252 to 65 mya. This era is further divided into the Triassic, Jurassic, and Cretaceous periods. The dinosaurs died out at the end of the Cretaceous Period, around 65 mya. The earliest hominids (members of the human family) appeared around 4.4 mya.



Eoraptor

First dinosaurs

Dinosaurs first evolved in the Triassic Period. Early dinosaurs were small, two-legged animals, such as *Eoraptor*.

Triassic

Jurassic

MESOZOIC ERA


252 mya

200 mya

Jurassic giants


Dinosaurs continued to evolve in the Jurassic, and became the dominant life-forms on land. Some grew to huge sizes, such as the sauropods, while others, such as the theropod *Archaeopteryx*, took to the air.

Mamenchisaurus,
a sauropod



Primates emerge

The era after the Mesozoic is called the Cenozoic Era. It contains the Paleogene Period, in which primates, the mammalian ancestors of humans, first appeared.



Eosimias, an early primate

Coming of humans

Many modern mammals, including horses, camels, and cows, evolved in the Neogene period. Hominids—the ancestors of humans—appeared in Africa and spread across the world.



Cretaceous

Paleogene

Neogene

CENOZOIC ERA

65 mya

23 mya


Present day

We live in what is called the Quaternary Period, which started 2.6 mya and continues to the present day.

Dinosaurs die out

The Cretaceous Period saw the emergence of new types of dinosaur, including the horned ceratopsians and armored ankylosaurs. Around 65 mya, an asteroid or comet collided with the Earth, killing off the dinosaurs and marking the end of the Mesozoic Era.





The Triassic

The Triassic Period lasted from 252 mya to 200 mya. During this period, the Earth's continents were joined together as a single landmass called Pangaea. Just before the Triassic, a mass extinction had wiped out a vast number of life-forms, including most land animals. The empty habitats began to be filled by a range of reptiles and eventually, the first dinosaurs. Mammals also appeared at this time.


Pleuromeia

A changing world

Deserts spanned the globe during the Triassic, and most plants grew near coasts and in river valleys, where the soil was moist. Ferns and treelike plants, such as *Pleuromeia*, flourished during this time.



Fern frond



Early fliers


Some reptiles took to the air for the first time during the Triassic. These were the pterosaurs, which were close relatives of the dinosaurs.



Eudimorphodon,
an early pterosaur

Plant-eaters

Large animals that roamed the land during the Triassic included the rhynchosaurs, a group of piglike, plant-eating reptiles.



Hyperodapedon,
a rhynchosaur

Dinosaurs

The earliest known dinosaurs, such as *Coelophysis*, appeared in the Triassic, some 230 mya. Most dinosaurs of this time were small, about the height of a man, and greatly outnumbered by other reptiles.



Coelophysis



The Jurassic



The Jurassic Period started around 200 mya, and lasted for 54 million years. Around 175 mya, Pangaea began breaking up into two large continents—Laurasia and Gondwana, as seen on the left.


Scientists believe that a mass extinction at the start of the Jurassic wiped out most of the non-dinosaur reptiles of the Triassic. This reduced the competition for food and allowed dinosaurs to flourish.

Sea monsters

During the Mesozoic Era, the seas were ruled by giant reptiles. In the Jurassic, these included the dolphinlike ichthyosaurs and the lizardlike plesiosaurs.



Ichthyosaurus, an ichthyosaur




Williamsonia

A new environment

The Triassic deserts transformed into lush green forests, filled with plants that had developed in the Triassic. These included coniferous trees, such as the monkey puzzle—a plant still present today—and *Williamsonia*, a small tree with palmlike fronds.

Leaves of
a monkey
puzzle tree





Pterodactylus


Flying aces

The pterosaurs of the Triassic evolved into the more efficient fliers of the Jurassic, such as *Pterodactylus*. This pterosaur had longer wings and a shorter tail, making it more agile in the air.

The age of giants

The colossal sauropods were the largest creatures ever to walk on the Earth, some of them reaching up to 59 ft (18 m) in height. These giants thundered through the forests in herds and they were often preyed upon by the largest theropods—ferocious meat-eating dinosaurs.

Brachiosaurus,
a sauropod



The Cretaceous



Lasting from 145 mya until 65 mya, the Cretaceous Period saw a number of changes across the globe. Laurasia and Gondwana broke up and began taking on the positions and shapes of continents today. The sauropods began to decline in number, while other plant-eaters, such as the duck-billed dinosaurs and ceratopsians, flourished. Flowering plants also began appearing in this period.

Adding color

In the Early Cretaceous, lush green forests of conifers and ferns still covered most of the land. One plant of the time was the tree-fern *Tempskya*. Instead of a trunk, it had a number of shoots joined together.

Eventually, the landscape began to change. Flowering plants, such as magnolia, appeared, adding color to the vegetation.



Magnolia
flower

Tempskya





Liaoxiornis,
an early bird

Feathered fliers

Although the first birds evolved in the Jurassic, those that appeared in the Cretaceous were more varied. Some had toothless beaks and other features similar to today's birds.



Mosasaurus,
a mosasaur

Ambush predators

During the Cretaceous, the oceans were ruled by gigantic predators called mosasaurs. Their powerful tails made them good swimmers, but these giants preferred to lie in wait before pouncing on their prey, rather than chasing it down.



Zalambdalestes

Little scavengers

Mammals lived throughout the age of the dinosaurs, but remained small, feeding on plants, small animals, and eggs.

Jungle grazer

Ceratopsians, along with duck-billed dinosaurs, such as *Maiasaura*, became widespread in the Cretaceous. *Triceratops* was a large plant-eating ceratopsian that lived in herds, probably feeding on flowering plants.



Triceratops

Dinosaur ancestors

Dinosaurs, like all land animals with backbones, evolved from fish. The fins of the fish developed into limbs that helped in walking on land, while lungs developed and allowed breathing. One group of these animals evolved into the reptiles, and some reptiles evolved into dinosaurs.




Panderichthys

Fishy bones

Lobe-finned fish, such as *Panderichthys*, were the ancestors of all four-legged backboned animals, or tetrapods. Their fins sprouted from fleshy structures, or lobes, strengthened by bones like those in our limbs.

First steps

Acanthostega was one of the earliest tetrapods. It had a fishlike tail fin that pushed it through the water, and four limbs, each with eight toes, which allowed it to walk.



Paddlelike tail fin

Leg with eight toelike digits


Acanthostega

Protecting the eggs

The eggs of early tetrapods needed to be laid in water. Over time, some tetrapods, such as *Westlothiana*, evolved eggs with a waterproof membrane. This allowed them to lay their eggs on land, and the eggs didn't dry out. These animals began breeding on land, evolving into reptiles, dinosaurs, and mammals.



Westlothiana



Bony plates
along back

Sideways
facing eyes


Sprawling
front limbs

Parasuchus

Dinosaur cousins

Dinosaurs are part of a group of reptiles called archosaurs. Modern crocodiles also belong to this group and evolved from early archosaurs, such as *Parasuchus*. The early archosaurs moved with their knees bent, giving them a sprawling walk.

Postosuchus, an advanced archosaur



On two legs


As archosaurs evolved, some began to stand upright, and unlike the sprawling legs of crocodiles, their legs supported the body high above the ground, allowing agile and efficient running. Scientists believe that dinosaurs may have evolved from these kinds of upright archosaurs.

Types of dinosaur

More than 1,000 species of dinosaur lived in the Mesozoic. Enormous sauropods grazed on the very tops of trees, while packs of meat-eating theropods chased down their prey and ripped it apart with sharp teeth and claws. Some dinosaurs had armored skins, while others had horns or spikes to protect themselves.

Huge!

Some dinosaurs were the largest animals ever to walk on the Earth, but no one is sure why they grew so big. Some scientists suggest size developed as a defense against predators. Others propose that an abundance of food led to these giants.



Barosaurus, a sauropod, length 92 ft (28 m)

Muttaborrasaurus, an ornithomimid, length 26 ft (8 m)

Ankylosaurus, an ankylosaur, length 20 ft (6 m)



Dinosaurs

Saurischians
Ornithischians

Family tree

Early dinosaurs evolved into two main groups—the ornithischians and the saurischians. Saurischians include the meat-eating theropods and the long-necked sauropodomorphs—a large group containing the sauropods and prosauropods. Ornithischians also branched into different types. The armored ankylosaurs and stegosaurs broke off first, with the three-toed ornithopods and frilled ceratopsians and pachycephalosaurs appearing later.

Theropods



Sauropodomorphs



Ankylosaurs



Stegosaurs




Ornithopods

Pachycephalosaurs



Ceratopsians




Styracosaurus, a ceratopsian, length 17 ft (5.2 m)

Alxasaurus, a theropod, length 13 ft (4 m)

Modern man, average height 6 ft (1.8 m)

Dinosaurs to birds

We now know that birds evolved from theropod dinosaurs. However, the features that define modern birds—flight feathers, short tails, and a curved breastbone with strong flight muscles that allow them to fly—were not present in their theropod ancestors. It took millions of years for these features to appear.



Feathered find


When it was first discovered in 1996, *Sinosauropteryx* caused a huge sensation among scientists. It had a soft, fuzzy covering that resembled feathers. Scientists realized that dinosaurs evolved feathers before taking to the air.

Using feathers

The first feathers were not used for flight. Some feathers helped keep the theropods warm, while others were used for display. We know this because of the well-preserved fossils of *Sinornithosaurus*, which show remains of feathers of various kinds.



A complete fossil of *Sinornithosaurus*



Micro flier

One of the smallest dinosaurs, *Microraptor* was slightly bigger than a pigeon. This tiny theropod had asymmetrical feathers on its arms and legs that generated lift, allowing it to glide from tree to tree.

First bird

Once considered the first true bird, *Archaeopteryx* is one of the earliest known theropods capable of weakly powered flight. Its asymmetrical feathers also allowed it to glide. This Jurassic dinosaur, however, still had a long reptilelike tail and claws on its wings.



Modern bird

Iberomesornis was a finch-sized creature that lived in the Cretaceous and was one of the many birds that gave rise to modern birds. It had a short tail with tail feathers, and a curved breastbone, but lacked the strong flight muscles of modern birds.



EVOLUTION OF FEATHERS



Hollow hairlike filaments were the first feathers.



Tufted barbs joined at the base developed from the single filaments.



Barbs connected to a central axis developed next.




Asymmetrical feathers evolved. These could provide lift for flying.

Dinosaurs die out


The dinosaurs dominated the Earth for more than 160 million years. Then, about 65 mya, they disappeared in a mass extinction of many life-forms. There are many theories about why dinosaurs died out. The strongest evidence shows that an asteroid or comet crashed into the Earth. At the same time, volcanoes erupted. Both events led to drastic climate change.

Death from the skies

Scientists know that an asteroid or comet, 6 miles (10 km) across, collided with the Earth at a very high speed. The impact threw up huge quantities of dust into the air, blocking out the Sun. Without the Sun's heat, temperatures across the Earth dropped rapidly, and most life-forms perished. This coincided with an increase in volcanic activity. Erupting volcanoes would have released large amounts of ash and toxic gases, destroying animal and plant life.



EVIDENCE



Deep impact Scientists know that the Chicxulub crater in the Yucatán Peninsula of Mexico was made by an asteroid or comet impact 65 mya. This artistic depiction shows how the impact crater might have looked from space. The crater was discovered in the 1990s and is more than 110 miles (180 km) across.



Volcanic activity This mass of rock is the Deccan Traps in India. It was formed by lava flow. One of the largest volcanic features on the Earth, it was built up by a series of volcanic eruptions 80 to 60 mya. It is estimated that the lava flow covered an area of 580,000 sq miles (1.5 million sq km)—half the size of modern India.



Phenacodus,
a mammal that
lived in the
Paleogene

Survivors

The survivors of the mass extinction included birds and the small, shrewlike mammals of the Cretaceous. With the theropods dead, there were no major predators left, and these mammals flourished, growing larger and becoming widespread.


How fossils form

Nearly everything we know about dinosaurs we learn from fossils. Fossils are the remains or traces of plants and animals that have actually turned to rock over millions of years. The rock usually forms as mineral-rich water trickles into the tiny spaces inside bones, shells, or tree trunks.

Fossilized


An animal only turns into a fossil when it dies if a few lucky things happen together. First, it must be buried quickly—perhaps by wind-blown sand or river mud. Over the millennia, the animal's skeleton changes into rock, or leaves an impression on the surrounding rock.

Decaying body




1. **A fish sinks** to the riverbed when it dies. Its soft parts might rot or be eaten.

Sediments such as sand or mud build over skeleton




2. **Minerals** in the water may seep into the tiny spaces in the bone and crystallize.

Layers of sediment squash fossil



3. **With time**, heat, and pressure, the mineral crystals in the bones turn into rock.

Fossilized fish skeleton



Rock exposed due to erosion by wind and water

4. **Over millions of years**, the layers of rock above may be worn away, leaving the skeleton at the Earth's surface—ready for fossil hunters to discover.



Mineral invasion

Ammonites were squidlike creatures that lived inside a coiled shell. They existed at the same time as the dinosaurs. The spaces inside this ammonite's fossil shell have been filled with a mineral called pyrite, also known as fool's gold. The shell itself has been transformed into a darker mineral.

Soft parts preserved

Usually, only hard parts of an animal, such as shell and bone, are fossilized, because soft parts are eaten or rot away before they are buried. In rare cases of a very quick burial, skin and other soft parts can be preserved as rock.



Skin of *Xiphactinus*, a prehistoric fish

Trace fossils

Fossils are not always the remains of animal and plant bodies. They can be marks and signs they left behind, such as eggs, droppings, and footprints. These signs are called trace fossils.



Model of *Allosaurus* and *Apatosaurus* footprints

Fossil bones

Whole skeletons are rarely discovered intact. Most fossils are single teeth, bones, or partial skeletons. Paleontologists must use this scant evidence to piece together the missing parts. Sometimes, fossil hunters are lucky enough to find whole skeletons, some of which are even still “articulated” (all bones in place).

Iguanodon hand fossil

A thumb spike can be clearly seen in this nicely articulated *Iguanodon* fossil, but scientists have not always had such good evidence of this animal. *Iguanodon* was one of the first dinosaurs to be discovered. In 1820, English doctor Gideon Mantell found fossil teeth in a quarry and realized that they belonged to a giant prehistoric animal. Soon afterward, Mantell and others found more nonarticulated *Iguanodon* bones. Limited fossil evidence was easy to misinterpret. In piecing together the creature, Mantell mistakenly placed the thumb spike as a horn on its nose!

DATE 135–125 mya
(Early Cretaceous)

SIZE 10 in (25 cm) long

FOSIL LOCATION England

ANIMAL GROUP Dinosaurs



Gryposaurus skeleton

Gryposaurus was a hadrosaur (see pp. 92–93). This skeleton was found in Alberta, Canada, at a site called the Dinosaur Park Formation. It is partially embedded in rock and partially mounted. Scientists mount such complete fossil specimens based on their understanding of how the creature stood and moved. Intact fossils such as this show that strong tendons held up the tail, preventing it from sagging. Earlier reconstructions of these creatures pictured them standing upright.



DATE 83–75 mya (Late Cretaceous)

SIZE 29½ ft (9 m) long

FOSSIL LOCATION North America

ANIMAL GROUP Dinosaurs

Baryonyx claw fossil

Scientists don't know for certain if this *Baryonyx* claw came from its index finger or thumb. Fossils of single claws found separated from the body can be misleading. The claw had a groove where a sheath of horn was attached.

DATE 125 mya (Early Cretaceous)

SIZE 14 in (35 cm)
from tip to base

FOSSIL LOCATION
England

ANIMAL GROUP
Dinosaurs



Gallimimus skull fossil

Gallimimus had a birdlike skull with a long toothless beak. It had wide eye sockets with eyes facing sideways. Each eyeball was supported by a ring of bony plates.

DATE 75–65 mya (Late Cretaceous)

SIZE 12 in (30 cm) long

FOSSIL LOCATION Mongolia

ANIMAL GROUP Dinosaurs



Exceptional fossils

The hard parts of an animal's body tend to fossilize the best. In exceptional cases, however, an animal might be buried rapidly with no chance for the soft body parts to rot. The resulting fossils include those of skin and feathers, and even internal organs.

Edmontosaurus skin fossil



Mud filled every crease in the skin of *Edmontosaurus* to create this fine fossil, called a cast, which shows what the scales on the skin looked like.



DATE 75–65 mya (Late Cretaceous)

SIZE 8 in (20 cm) across

FOSSIL LOCATION United States

ANIMAL GROUP Dinosaurs

Polacanthus skin fossil



Large, spiked lump

This *Polacanthus* fossil has traces of the animal's knobby skin. Mud covered the animal before its body had decayed and a mold (impression) of its skin was filled by the mud. This turned to rock, preserving the spiked lumps on the skin of this ankylosaur.

DATE 130 mya (Early Cretaceous)

SIZE 6 in (15 cm) across

FOSSIL LOCATION England

ANIMAL GROUP Dinosaurs

Sinornithosaurus fossil

In 2001, paleontologists discovered this complete fossil of *Sinornithosaurus* in China. Impressions of primitive feathers line the bones of this creature. This dinosaur was a ground-dwelling animal and probably died on a riverbed with a fish in its claws. It was buried so quickly that its feathers were intact and left their shapes on the surrounding mud before rotting away. This discovery helped scientists understand that not all kinds of feather were used for flying. Some kinds helped keep the feathered theropods warm.



Feather impression

Fossil fish

DATE 130–125 mya (Early Cretaceous)

SIZE 3¼ ft (1 m) long

FOSSIL LOCATION China

ANIMAL GROUP Dinosaurs



Sinornithosaurus means “Chinese bird lizard,” but this creature was not a true bird and couldn’t fly.

Trace fossils

Sometimes an animal leaves behind a hint, or trace, of its presence. The remains of such traces of prehistoric creatures and their activities, preserved in rock, are called trace fossils. These include footprints, bite marks, droppings, and eggs.

Ichthyosaur coprolite



Coprolites are the fossilized dung of prehistoric creatures. Scientists study these fossil droppings to understand more about the diets of the animals. This is the coprolite of a marine reptile called an ichthyosaur (see pp. 124–25). Undigested scraps of bone and shell from its last meal can reveal the kinds of prey it fed on.

DATE 190 mya (Early Jurassic)

SIZE 3¼ in (8 cm) long

FOSSIL LOCATION England

ANIMAL GROUP

Ichthyosaurs



Apatosaurus egg fossil



This is a fossilized egg of the sauropod *Apatosaurus* (see pp. 60–61). The eggs of sauropods had a thick shell that protected them from breakage. The eggs seem small in comparison to the sizes of these giants, but larger eggs would have needed shells so thick that hatchlings could not have broken out.

DATE 154–150 mya (Late Jurassic)

SIZE 5 in (13 cm) across

FOSSIL LOCATION United States

ANIMAL GROUP Dinosaurs

Oviraptor egg and embryo fossils

These are the fossilized bones of an embryo of the theropod *Oviraptor*. They lie within the fossil remains of an egg shell and were found in a fossilized nest in the Gobi Desert. Finding the delicate bones of an embryo like this helps scientists figure out which dinosaur laid the eggs.

DATE 75 mya (Late Cretaceous)

SIZE 7 in (18 cm) long

FOSSIL LOCATION Mongolia

ANIMAL GROUP Dinosaurs



Iguanodon footprint

A young *Iguanodon* left this footprint in the Early Cretaceous. Sometimes footprints get preserved in layers of mud, which hardens to rock, in turn preserving the footprints as fossils. The shapes and sizes of these prints and the gaps between them allow scientists to identify the kinds of dinosaur that made the prints, and also to figure out the sizes of the dinosaurs and how fast they walked or ran. Based on the length of this footprint, scientists determined that it was made by an *Iguanodon* weighing roughly half a ton.

DATE 135–125 mya (Early Cretaceous)

SIZE 11½ in (29 cm) long

FOSSIL LOCATION England

ANIMAL GROUP Dinosaurs




Reconstruction

Reconstructing dinosaurs requires a lot of research. After excavating fossils, scientists study them and compare them to modern reptiles to understand how the bones in a dinosaur skeleton connected to each other and to the muscles that moved a dinosaur's body.


Mounting dinosaurs

When arranging a dinosaur skeleton, scientists not only have to make sure of how the bones connect, but they also have to fix the dinosaur's pose. Sometimes, new research leads to resetting of the posture. For example, it was once thought that sauropods dragged their tails on the ground, but we now know that they held their tails aloft.



Let's move


Dinosaurs can be re-created as moving 3-D digital models using CGI, or computer-generated imagery. This is how the *Corythosaurus* image (p. 93) was built.




1. Scientists and computer programmers study the dinosaur's bones and make a basic model from geometrical shapes on a computer. This is called a wireframe.




2. A computer program divides the geometrical figures into millions of smaller units. An artist can then shape each unit, refining the dinosaur's shape.



3. New discoveries about the dinosaur, such as the size and shape of its crest, are added to the sculpture. This helps create an accurate model of the dinosaur.



4. Adding color to the dinosaur involves a mix of artistic creativity and scientific insight. Fossil dinosaur skin may feature intact microscopic pigment capsules, the shapes of which help scientists deduce the dinosaur's color.



5. A rigger (a specialized programmer with expertise in anatomy) tells the computer how the different parts of the body would have moved.



6. Scientists tell the artist what kind of environment the dinosaur lived in, allowing the artist to create realistic backgrounds of the dinosaur's habitat.



Dinosaurs

Dinosaurs were the dominant land animals for more than 160 million years and ranged from small animals no bigger than pigeons to lumbering giants the size of a truck. As scientists have studied remains of these ancient reptiles, they have found evidence of the original colors of these creatures in their feathers. Many dinosaurs had striped or bright feathers, and probably also multicolored skin, feathers, frills, and head crests, which made them look more attractive or fierce.



JURASSIC FOREST

Thick, lush forest covered the land during the Jurassic, with trees and ferns providing food and shelter for the dinosaurs.

The first dinosaurs

Dinosaurs evolved in the Triassic Period. The earliest ones were agile animals that walked on their hind legs and had sharp teeth and claws. Many of them were omnivores that fed on a variety of foods. They gradually evolved into specialized plant-eating and meat-eating dinosaurs.

Herrerasaurus

her-air-ah-SORE-us



Slender, flexible neck was typical of most early dinosaurs

Herrerasaurus is one of the most primitive dinosaurs and lived in a world dominated by non-dinosaur reptiles. Its short thighs and long feet made it a fast runner, like most of the earliest dinosaurs, and it could probably have easily outrun its prey.

DATE 231 mya (Late Triassic)

SIZE 10–20 ft (3–6 m) long

FOSSIL LOCATION Argentina

HABITAT Forests

DIET Animals

Eoraptor

ee-oh-RAP-ter




One of the earliest saurischians, *Eoraptor* was the size of a fox. It had the sawlike teeth of a hunter. Its eyes faced toward the side.

DATE 231 mya (Late Triassic)

SIZE 3¼ ft (1 m) long

FOSSIL LOCATION Argentina

HABITAT Forests

DIET Lizards, small reptiles, and plants
**Eocursor**

ee-oh-CUR-ser




Eocursor was an early ornithischian. This omnivore could run quickly on its hind legs. Its hands were armed with sharp claws that probably helped in catching small animals.

DATE 210 mya
(Late Triassic)

SIZE 3¼ ft (1 m) long

FOSSIL LOCATION
South Africa

HABITAT Moist forests

DIET Plants, small mammals, and reptiles
**Gojirasaurus**

go-JEER-a-SORE-us




The movie monster Godzilla, known as *Gojira* in Japanese, inspired the name of this meat-eater. It prowled across the arid regions of what is now North America, and it was the top predator of the American Southwest.

DATE 210 mya (Late Triassic)

SIZE 16½–23 ft (5–7 m) long

FOSSIL LOCATION United States

HABITAT Scrublands

DIET Animals




FOCUS ON...

JAW DIVERSITY

The jaws of theropods were suited to catching and eating in different ways.



▲ *Baryonyx* had narrow jaws with pointed teeth, good for catching fish.



▲ *Deinonychus* had pointed jaws, good for tearing out pieces of flesh.




▲ *Tyrannosaurus* had big, U-shaped jaws, which tore out large chunks of meat.

Theropods

A varied group, theropods included all predatory dinosaurs, some omnivores, and probably some plant-eaters. Some predatory theropods were tiny, some were birdlike, and others were giant top predators. They killed prey with bladelike teeth and sharp claws.

Coelophysis

SEE-low-FYE-sis



This lightweight hunter had more than 100 small, curved teeth in its narrow jaws. Its long, flexible neck helped it to dart its head forward when catching small prey.


DATE 208–200 mya (Late Triassic)

SIZE 9 ft (2.8 m) long

FOSSIL LOCATION United States

HABITAT Desert plains

DIET Lizards and fish



Ceratosaurus

ser-AT-oh-SORE-us



Ceratosaurus's name means "horned lizard" and refers to the rounded horn above its nostrils. It also had a horn in front of each eye. A row of low spines ran down its neck, back, and tail—an uncommon feature for a theropod.

DATE 150–144 mya (Late Jurassic)

SIZE 20 ft (6 m) long

FOSSIL LOCATIONS


United States and Portugal

HABITAT Forests

DIET Dinosaurs and other reptiles

Liliensternus

LIL-ee-en-STERN-us



In its time, *Liliensternus* was the largest predator on land. Its long hind legs allowed it to run fast after its prosauropod prey. This theropod probably stalked its victims before attacking.

DATE 210 mya (Late Triassic)

SIZE 16½–20 ft (5–6 m) long

FOSSIL LOCATION Germany

HABITAT Forests


DIET Dinosaurs

Dilophosaurus

di-LOAF-oh-SORE-us



This animal is best-known for the two parallel, platelike crests on its snout. Scientists believe these were used in a display for attracting mates.



DATE 201–189 mya
(Early Jurassic)

SIZE 20 ft (6 m) long

FOSSIL LOCATION US

HABITAT Riverbanks

DIET Small animals and fish

Monolophosaurus

mono-LOAF-oh-SORE-us



With a thick crest on its head, *Monolophosaurus* had one of the biggest and oddest of skulls. The hollow crest may have helped to produce noises that warned off rivals.



DATE 180–159 mya
(Middle Jurassic)

SIZE 20 ft (6 m) long

FOSSIL LOCATION
China

HABITAT Forests

DIET Dinosaurs



Cryolophosaurus

CRY-oh-LOAF-oh-SORE-us

The largest known theropod from the Early Jurassic, *Cryolophosaurus* had slender arms and long legs. It was also the first theropod to be discovered in Antarctica. The crest on its head was unusual in shape, curving upward and forward over the skull.

DATE 190–185 mya (Early Jurassic)

SIZE 21½ ft (6.5 m) long

FOSSIL LOCATION Antarctica

HABITAT Open plains

DIET Dinosaurs



People nicknamed this dinosaur *Elvisaurus* because its head crest reminded them of Elvis Presley's hair.

Baryonyx

bah-ree-ON-ix

A fish-eating theropod, *Baryonyx* had an unusual curved claw on its thumb or index finger. It used the claws as hooks to kill its prey, much like bears do today.


DATE 125 mya (Early Cretaceous)

SIZE 29½ ft (9 m) long

FOSSIL LOCATIONS British Isles, Spain, and Portugal

HABITAT Riverbanks

DIET Fish and dinosaurs



Suchomimus

soo-ko-MIME-us

A close relative of *Baryonyx*, this predator had a long, crocodilelike snout and a slim jaw with more than 100 teeth that pointed backward. *Suchomimus* used its teeth and long arms to hold slippery prey.

Teeth at tip of snout were longer than those at the back

DATE 112 mya (Early Cretaceous)


SIZE 29½ ft (9 m) long

FOSSIL LOCATION Africa

HABITAT Mangrove swamps

DIET Fish and possibly other animals

Crocodilelike jaw



Spinosaurus

SPINE-oh-SORE-us

Spinosaurus is one of the longest theropods known. It had a sail-like structure running down its back, supported by spines made of bone. This gave it the name *Spinosaurus*, meaning “spine lizard.” Scientists believe that this predator may have used its sail to control its body temperature.


DATE 97 mya (Late Cretaceous)

SIZE 59 ft (18 m) long

FOSSIL LOCATIONS Morocco, Libya, and Egypt

HABITAT Tropical swamps


DIET Fish and dinosaurs



Raised on bones 6½ ft (2 m) long, the sail of *Spinosaurus* was an impressive sight.

Allosaurus

al-oh-SORE-us



This fierce predator had a massive skull, powerful jaws, and a long tail. Sturdy bones supported its jaw and bladelike teeth. The skull enabled *Allosaurus* to chop flesh rapidly, unlike the slow, bone-crushing skull of *Tyrannosaurus*.

DATE 150 mya (Late Jurassic)

SIZE 39 ft (12 m) long

FOSSIL LOCATIONS

US and Portugal

HABITAT Open plains

DIET Large
plant-eating
dinosaurs

Carcharodontosaurus

CAR-ka-roe-DON-toe-SORE-us

Twice the weight of an elephant, this enormous theropod used its massive jaws equipped with sawlike teeth to kill prey. The scientists who first discovered it found its teeth similar to those of the great white shark, *Carcharodon*, and named it *Carcharodontosaurus*.

DATE 100 mya (Early Cretaceous)


SIZE 46 ft (14 m) long

FOSSIL LOCATIONS

Morocco, Tunisia, and Egypt

HABITAT Floodplains and mangroves

DIET Large plant-eating dinosaurs



Giganotosaurus

GEEG-ah-NOTE-ih-SORE-us

Similar in size to *Tyrannosaurus*, *Giganotosaurus* was as heavy as 125 people. Despite its size, it could probably run at 30 mph (50 kph) when chasing its prey.


DATE 112–90 mya (Early Cretaceous)

SIZE 43 ft (13 m) long

FOSSIL LOCATION Argentina

HABITAT Warm swamps

DIET Large dinosaurs



Sinraptor

SIN-rap-ter

A close relative of *Allosaurus*, *Sinraptor* was a formidable hunter. Tooth marks on a skull suggest that it might have fought with its own kind.

DATE 169–142 mya (Middle–Late Jurassic)

SIZE 24½ ft (7.5 m) long

FOSSIL LOCATION China

HABITAT Forests

DIET Large plant-eating dinosaurs



Powerful jaws lined with
bone-crunching
blades made *Tarbosaurus* the
most fearsome predator on
the plains of China



TARBOSAURUS

Tarbosaurus was a close relative of *Tyrannosaurus*, and both of them belong to a family of theropods called tyrannosaurids. *Tarbosaurus* preyed on smaller dinosaurs, such as *Barsboldia*, which was a hadrosaur.




Albertosaurus

al-BERT-oh-SORE-us



This lightly built predator had slender hind limbs and small forelimbs and may have been a swift runner. Some scientists believe that *Albertosaurus* lived and hunted in packs.



DATE 75 mya
(Late Cretaceous)

SIZE 29½ ft (9 m) long

FOSSIL LOCATION Canada

HABITAT Forests

DIET Dinosaurs

Compsognathus

COMP-sog-NAITH-us



Running on the tips of its toes, this chicken-sized predator could easily outpace fast-moving prey. Its long tail helped in balancing and making sharp turns while running.



DATE 150 mya (Late Jurassic)

SIZE 4¼ ft (1.3 m) long

FOSSIL LOCATIONS
Germany and France

HABITAT Scrublands
and marshes

DIET Lizards, mammals,
and small dinosaurs

Tyrannosaurus

TIE-ran-oh-SORE-us

Tyrannosaurus was as long as a bus and twice as heavy as an elephant. This fearsome predator had pointed teeth that could tear skin and muscle, and even crush bone. Three-toed feet allowed this heavy animal to plod along at a steady trot. Its tiny arms, with two claws on each, were probably used to hold the prey during feeding.

DATE 70–65 mya (Late Cretaceous)

SIZE 39 ft (12 m) long

FOSSIL LOCATION North America

HABITAT Forests and swamps

DIET Large dinosaurs





Although shown in the film *Jurassic Park*, *Tyrannosaurus* actually lived in the Cretaceous Period.




Guanlong

GWON-long



Guanlong means “crowned dragon,” the name reflecting the striking crest on the theropod’s head. It was a close relative of early feathered dinosaurs and had a coat of fuzzy feathers itself.



DATE 160 mya (Late Jurassic)

SIZE 8¼ ft (2.5 m) long

FOSSIL LOCATION China

HABITAT Forests

DIET Dinosaurs and other animals

Proceratosaurus

PRO-seh-RAT-oh-SORE-us



The only fossil of this crested dinosaur is this skull found in 1910. It is thought to be a small dinosaur and a close relative of *Guanlong*.

DATE 175 mya (Middle Jurassic)

SIZE 6½ ft (2 m) long

FOSSIL LOCATION British Isles

HABITAT Forests

DIET Dinosaurs and other animals




Gallimimus

GAL-ih-MIME-us



Although *Gallimimus* means “chicken-mimic,” this dinosaur was a lot heavier than a chicken. It was three times taller than a man and weighed around 1,000 lb (450 kg).



DATE 75–65 mya (Late Cretaceous)

SIZE 20 ft (6 m) long


FOSSIL LOCATION Canada

HABITAT Desert plains

DIET Leaves, seeds, insects, and small animals

Ornithomimus


OR-nith-oh-MIME-us



Ornithomimus belongs to a family of dinosaurs known as ornithomimids, or ostrich dinosaurs. This theropod was a fast runner, with a long, stiff tail that allowed it to make sudden turns while sprinting. Compared to other dinosaurs, it had a fairly large brain, but was far less intelligent than an ostrich.

Ornitholestes

OR-nith-o-LESS-teez



Long, powerful grasping fingers

A small and lightly built body made *Ornitholestes* a swift and efficient predator. It had long front teeth with flattened tips that helped it catch prey.


DATE 156–145 mya (Late Jurassic)

SIZE 6½ ft (2 m) long

FOSSIL LOCATION United States

HABITAT Forests

DIET Small animals, such as insects, lizards, and frogs



Long, powerful fingers with claws allowed for grasping plants and prey



DATE 75–65 mya (Late Cretaceous)

SIZE 10 ft (3 m) long

FOSSIL LOCATIONS US and Canada

HABITAT Swamps and forests

DIET Plants, seeds, and small animals

Caudipteryx

caw-DIP-ter-ix



Although its body was covered in feathers, *Caudipteryx* was unable to fly.

Scientists believe it used its feathers for display and to keep itself warm.



DATE 130–120 mya (Early Cretaceous)

SIZE 3¼ ft (1 m) long

FOSSIL LOCATION China

HABITAT Lake sides and riverbeds

DIET Plants, seeds, and small animals

Citipati

SIH-tee-PAH-tee



Citipati had a distinctive crest on its head that was made of horny keratin. In addition to plants, this theropod probably also fed on eggs and baby dinosaurs, ripping them apart with its beak, just as eagles do today.

DATE 75 mya (Late Cretaceous)

SIZE 10 ft (3 m) long

FOSSIL LOCATION Mongolia

HABITAT Open plains

DIET Plants and animals





Troodon had the largest brain of all dinosaurs, relative to its body size.

Microraptor

my-CROW-rap-ter

One of the smallest dinosaurs, *Microraptor* had long, birdlike feathers on its arms and legs. However, unlike birds, which flap their feathered wings and fly, the wings of *Microraptor* were not big enough to support the animal's weight and so it could only glide from branch to branch. It probably did so to search for prey and to escape predators.



DATE 130–125 mya (Early Cretaceous)


SIZE 3¼ ft (1 m) long

FOSSIL LOCATION China

HABITAT Forests

DIET Small mammals, lizards, and insects





Troodon

TROH-o-don



An unusually large brain for a dinosaur and sharp, forward-facing eyes made *Troodon* an effective hunter. Because its eyes faced forward, it could judge the distance to its prey before pouncing. This predator had strong, slender legs that allowed it to outrun most small animals.

DATE 74–65 mya (Late Cretaceous)

SIZE 10 ft (3 m) long

FOSSIL LOCATION North America

HABITAT Forests

DIET Small animals and possibly plants

Velociraptor

vel-OSS-a-rap-ter



About the size of a wolf, this theropod used its long clawed arms to grapple and bring down prey. It is one of the best-known of all dinosaurs, having been shown in films such as *Jurassic Park*.



DATE 85 mya (Late Cretaceous)

SIZE 6½ ft (2 m) long

FOSSIL LOCATION Mongolia

HABITAT Scrublands and deserts

DIET Lizards, mammals, and small dinosaurs

Deinonychus

dye-NON-ee-cuss



Famous for its large toe claws, *Deinonychus* was a fierce predator. Some experts think that the sickle-shaped claws may have been used to slash the throat or belly of prey.



DATE 115–108 mya (Early Cretaceous)

SIZE 10 ft (3 m) long

FOSSIL LOCATION US

HABITAT Swamps and forests

DIET Small dinosaurs



Many *Citipati* fossils have been found crouching over eggs in nests, seeming to brood the eggs

just like a bird



CITIPATI

Citipati belongs to a family of theropods called oviraptorosaurs. These dinosaurs had parrotlike beaks and their bodies were covered with feathers. *Caudipteryx* and *Oviraptor* were also members of this family.

Early birds

Birds began as small, feathered, toothed dinosaurs with long, bony tails and small flight muscles. Over time, they evolved shorter tails, stronger muscles, and a lighter skeleton.



FOCUS ON... FEATURES

Early birds had many features that are absent in their modern cousins.

Archaeopteryx

ar-kee-OP-ter-ix



Archaeopteryx was capable of weakly powered flight. Weak flight muscles and a bony tail meant that it was a clumsy flier. It was long considered the earliest bird. Some scientists now think that a theropod called *Xiaotingia* may have been more closely related to early birds.

DATE 150 mya (Late Jurassic)

SIZE 12 in' (30 cm) long

FOSSIL LOCATION Germany

HABITAT Forests and lakes

DIET Insects and reptiles



▲ The fingers on the forelimbs had claws, which helped the early birds to climb.



▲ The tails of early birds were long and reptilelike, unlike the bony stump of modern birds.



▲ Early birds had teeth typical of theropods, unlike today's toothless birds.

Ichthyornis

ICK-thee-OR-niss



Like modern birds, this gull-sized seabird had a deep keelbone—an extension of the breastbone—that anchored its flight muscles. It also had a boxlike rib structure similar to modern birds. It was still a primitive bird, however, and had small, sharp teeth.



DATE 90–75 mya (Late Cretaceous)

SIZE 23½ in (60 cm) long

FOSSIL LOCATION United States

HABITAT Seashores

DIET Fish

Vegavis

VAY-gah-viss



Vegavis is distantly related to today's ducks and geese. The discovery of *Vegavis* shows that some of today's bird families had already evolved in the Cretaceous Period.

DATE 65 mya (Late Cretaceous)

SIZE 23½ in (60 cm) long

FOSSIL LOCATION Antarctica

HABITAT Seashores

DIET Water plants





FOCUS ON...

FEATURES

Some characteristics were common to all prosauropods.



▲ Prosauropods had a massive, clawed thumb used to rake in plants.



▲ Prosauropods had a high snout and long, slender jaws.



▲ Small, leaf-shaped teeth could easily slice through tough stems.

Prosauropods


These early plant-eating relatives of sauropods evolved from small meat-eating dinosaurs in the Triassic. Over time, they grew taller and heavier, with long necks and strong hind legs that let them reach high tree branches. They had hands with fingers and thumbs.

Efraasia

e-FRAHS-ee-a



Efraasia had a small head with a long neck. Its five-fingered hands had a large thumb claw. It may have walked on all fours to browse on leaves, but probably ran on its hind legs.



DATE 210 mya
(Late Triassic)

SIZE 20–23 ft
(6–7 m) long

FOSSIL LOCATION
Germany

HABITAT Dry plains

DIET Plants and
possibly animals

Thecodontosaurus

THEE-co-DON-toe-SORE-us

The unusual leaf-shaped, sawlike teeth of this prosauropod resembled those of a modern-day monitor lizard, but were attached to distinct sockets in its jaws. This inspired the name *Thecodontosaurus*, which means “socket-toothed lizard.”

DATE 225–208 mya (Late Triassic)

SIZE 6½ ft (2 m) long

FOSSIL LOCATION British Isles

HABITAT Island forests

DIET Plants and possibly animals



Anchisaurus

ankee-SORE-us

This early cousin of the sauropods had a shallow skull and a flexible spine. It had a narrow snout and pointed teeth in its upper jaw. *Anchisaurus* was probably an omnivore, feeding on small reptiles along with leaves.


DATE 190 mya (Early Jurassic)

SIZE 6½ ft (2 m) long

FOSSIL LOCATION United States

HABITAT Forests

DIET Leaves and small reptiles



Plateosaurus

PLATE-ee-oh-SORE-us



One of the best-known prosauropods, *Plateosaurus* could probably stand upright like a kangaroo, rearing on its hind limbs and stretching its legs to eat leaves from trees. Its sharp teeth could easily slice through tough leaf stems.

DATE 220–210 mya
(Late Triassic)

SIZE 26 ft (8 m) long

FOSSIL LOCATIONS Germany, Switzerland, Norway, and Greenland

HABITAT Open plains

DIET Plants



Lufengosaurus

loo-FENG-oh-SORE-us



This prosauropod had a massive claw on each of its thumbs that probably helped to grasp tree branches while feeding. *Lufengosaurus* used its widely spaced, bladelike teeth to rake leaves off branches.

DATE 200–180 mya (Early Jurassic)

SIZE 16½ ft (5 m) long

FOSSIL LOCATION China


HABITAT Forests

DIET Plants, including cycad and conifer leaves



Massospondylus

MASS-oh-SPON-dill-us



Fossils of *Massospondylus* show that it had a bulky body with a long tail. Five-fingered hands with massive thumb claws may have been used to tear off branches and stems. It may also have used its small, coarse teeth to chew on meat.

DATE 200–183 mya (Early Jurassic)

SIZE 13–20 ft (4–6 m) long

FOSSIL LOCATION South Africa

HABITAT Forests

DIET Plants and animals

Sauropods

The largest creatures ever to walk on the Earth, these lumbering giants had long necks and tails and pillarlike legs, but relatively small heads. They lived in herds and walked on all fours.



**FOCUS ON...
FOOTPRINTS**
Sauropod footprints have been found across the world.

Diplodocus

dip-LOD-oh-kuss

Probably one of the longest dinosaurs ever, *Diplodocus* had a tail that was as long as the rest of its body. Some scientists think that it raised its neck to browse on treetops, while others believe that it swung its head from side to side, feeding on shrubs.

DATE 150–145 mya (Late Jurassic)

SIZE 98½–110 ft (30–33.5 m) long

FOSSIL LOCATION United States

HABITAT Plains

DIET Plants

Tail bones become narrow at the tip



Diplodocus could move its tail at very high speed, producing a whiplike crack, which may have frightened off predators.





◀ In 1997, paleontologists discovered sauropod footprints in fossilized mud in a beach near Broome in Australia.



◀ The Purgatoire River track site in Colorado has more than 100 dinosaur trackways. Sauropod footprints can be seen here.

Apatosaurus

a-PAT-oh-SORE-us

Also known as *Brontosaurus*, this plant-eating giant was probably shorter than other sauropods and had thicker legs. *Apatosaurus* may have knocked down trees in search of food, much like elephants do today.

DATE 150 mya (Late Jurassic)

SIZE 75½ ft (23 m) long

FOSSIL LOCATION United States

HABITAT Forests

DIET Plants



Long, complex neck bones provide framework for supporting the head



A fully grown *Apatosaurus* could weigh as much as four elephants.

Barosaurus

BAH-roe-SORE-us

With a neck as long as 31 ft (9.5 m), this sauropod had an advantage over other dinosaurs when it came to reaching leaves right at the tops of trees. Its teeth were shaped like pegs, allowing it to rake leaves easily off the branches.

DATE 155–145 mya (Late Jurassic)

SIZE 92 ft (28 m) long

FOSSIL LOCATION United States

HABITAT Forests and plains

DIET Plants



Like today's cows, *Barosaurus* may have carried bacteria in its intestines that digested its food.



Amargasaurus

ah-MAR-gah-SORE-us

A double row of spines ran down *Amargasaurus's* neck, ending at its tail. There may have been a web of skin between the rows, forming a sail used for display.


DATE 130 mya (Early Cretaceous)

SIZE 36 ft (11 m) long

FOSSIL LOCATION Argentina

HABITAT Forests

DIET Plants





Dicraeosaurus

DIE-cray-oh-SORE-us



The bony spines running down the neck and back of *Dicraeosaurus* formed a ridgelike structure. It may have used the ridge for display, defense, or for controlling body temperature.

DATE 150 mya (Late Jurassic)


SIZE 39 ft (12 m) long

FOSSIL LOCATION

Tanzania

HABITAT Forests

DIET Plants



Vulcanodon

vul-KAN-o-don



This sauropod was named *Vulcanodon*, since its first fossils were found in rocks near volcanoes. Like other sauropods, it had short, elephantlike feet that were not suitable for running.


DATE 208–201 mya (Late Triassic)

SIZE 23 ft (7 m) long

FOSSIL LOCATION Zimbabwe

HABITAT Forests and plains

DIET Plants





Barosaurus could browse the treetops 49 ft (15 m) above the ground—the height of a **four-story building**




**BAROSAURUS**

A relative of the sauropod *Diplodocus*, *Barosaurus* had a very long neck. It had 15 cervical vertebrae (neck bones), some of which were more than 3¼ ft (1 m) long. This sauropod lived in herds, moving through the Jurassic forests in search of food.

Titanosaurus

tie-TAN-oh-SORE-us



Scientists believe that this dinosaur had bony studs on its back, much like its relative Saltasaurus

Titanosaurus is known only from fossil remains of limb bones. Many scientists believe that it had a typical sauropod body shape with a small head, short neck, and bulky body. Others, however, argue that the fossils belong to other dinosaurs, and *Titanosaurus* is not a separate species at all.

DATE 80–65 mya (Late Cretaceous)

SIZE 39–59 ft (12–18 m) long

FOSSIL LOCATIONS Asia, Europe, and Africa

HABITAT Forests and plains

DIET Plants

Saltasaurus

SALT-ah-SORE-us

Although smaller than many other sauropods, *Saltasaurus* had bony plates and studs running down its back, which defended it from large predators. Its neck was shorter than that of most sauropods and, unlike most of them, this animal lacked claws on its front feet.


DATE 80–65 mya (Late Cretaceous)


SIZE 39 ft (12 m) long

FOSSIL LOCATION Argentina

HABITAT Forests and open plains

DIET Plants





Argentinosaurus

ARE-jen-teen-oh-SORE-us

Argentinosaurus was one of the largest and heaviest land animals ever to walk on the Earth. It was longer than a tennis court and 20 times heavier than an elephant.


DATE 112–95 mya (Early Cretaceous)

SIZE 108–134 ft (33–41 m) long

FOSSIL LOCATION Argentina

HABITAT Forests and open plains

DIET Conifers



Mamenchisaurus

ma-MEN-chee-SORE-us

Mamenchisaurus had one of the longest necks of any known animal. The 19 long bones allowed it to move the neck freely and reach for food with great ease. It had a small, pointed head. It was named after a Chinese village, where its fossils were first found.


DATE 155–145 mya (Late Jurassic)

SIZE 85 ft (26 m) long

FOSSIL LOCATION China

HABITAT Riverbanks, forests, and open plains

DIET Trees and other plants



Brachiosaurus

BRACK-ee-oh-SORE-us

This sauropod used its long neck to feed on treetop leaves at heights greater than 49 ft (15 m), which is twice as high as any giraffe can reach. *Brachiosaurus* used its spoonlike teeth to snip off and eat an amazing 441 lb (200 kg) of leaves per day.

DATE 150–145 mya (Late Jurassic)

SIZE 75½ ft (23 m) long

FOSSIL LOCATION United States

HABITAT Forests and plains

DIET Leaves and twigs of conifers




Brachiosaurus weighed as much as 12 African elephants—an incredible 30–50 tons.



Camarasaurus

KAM-a-ra-SORE-us




Camarasaurus means “chambered lizard.” The dinosaur got this name because of the large air spaces inside some of the hollow bones connected to its lungs. These chambers helped reduce the animal’s body weight.

DATE 150–140 mya (Late Jurassic)

SIZE 59 ft (18 m) long

FOSSIL LOCATION United States

HABITAT Open plains

DIET Tree leaves
**Shunosaurus**

SHOO-noe-SORE-us



Scientists have found and studied several complete fossil skeletons of *Shunosaurus*. It had 25–26 teeth in each half of its lower jaw—more than any other sauropod.

Short, flexible neck

DATE 170–160 mya (Middle Jurassic)

SIZE 40 ft (12 m) long

FOSSIL LOCATION China

HABITAT Open plains

DIET Plants

Nemegtosaurus

nem-EGG-toe-SORE-us



A close relative of *Saltasaurus*, this sauropod is known only from a single fossil skull found in the 1970s. *Nemegtosaurus* was named after the Nemegt Basin in the Gobi Desert of Mongolia.


DATE 80–65 mya (Late Cretaceous)

SIZE 49 ft (15 m) long

FOSSIL LOCATION Mongolia

HABITAT Forests

DIET Plants





FOCUS ON...

BONY GROWTHS

The bony structures on stegosaurus' bodies had different functions.

Stegosaurus and relatives

Many dinosaurs evolved structures on their skin that helped them in defense against the predatory theropods. The stegosaurus, with rows of plates and spines along their backs, must have made a spectacular sight in the Jurassic forests.




▲ The plates on the back were probably used in courtship displays or perhaps to regulate body temperature by radiating heat away from the body.



▲ The long pointed spikes on the tail were used in defense against any predator that might attack from behind or the sides.

Scutellosaurus

SKOO-tell-oh-SORE-us



A relative of the stegosaurus, *Scutellosaurus* was an early ornithischian. This lightweight creature had hundreds of small bony scutes, or studs, on its skin.

DATE 196 mya (Early Jurassic)

SIZE 3¼ ft (1 m) long

FOSSIL LOCATION United States

HABITAT Forests

DIET Plants

Stegosaurus

STEG-oh-SORE-us



This animal was the largest of all the stegosaurs. It featured an alternating double row of flat, diamond-shaped plates running along its arched back. The plates were attached to the skin and were probably covered by keratin, the substance that forms horns and fingernails.

Stegosaurus's forelimbs were shorter than its hind limbs, so it walked with its hips higher than its shoulders. Its spine bones were tall and gave the animal a high, arched shape.

DATE 150–145 mya (Late Jurassic)

SIZE 29½ ft (9 m)

FOSSIL LOCATIONS

US and Portugal

HABITAT Forests

DIET Plants



This dinosaur was named *Stegosaurus*, meaning “roof lizard,” because its discoverer thought the plates on its back looked like roof tiles.



A predator would have broken
its teeth on the rows of


bony studs and spikes

running from *Scelidosaurus*'s
head to its tail



**SCELIDOSAURUS**

Scelidosaurus lived in the Early Jurassic and belonged to the same group of ancestral armored dinosaurs as *Scutellosaurus* (see p. 72). Its bony armor was covered in keratin—the material that forms nails and horns.




More than 900 *Kentrosaurus* bones were found in a single fossil site in Tanzania.

Huayangosaurus

HWAH-YANG-oh-SORE-us

Most stegosaurs have long hind limbs and short forelimbs, but all four of *Huayangosaurus*'s limbs were the same length. It was also different from later species as it had a shorter, broader snout, with teeth on the front of its upper jaw.



DATE 165 mya (Middle Jurassic)

SIZE 13 ft (4 m) long

FOSSIL LOCATION China

HABITAT River valleys

DIET Ferns, leaves, and cycad fruit

Kentrosaurus

KEN-troh-SORE-us



Seven pairs of bony plates lined the neck and back of this herbivore and may have been used for display. When attacked by a predator, *Kentrosaurus* probably lashed out its tail, using the long tail spikes to injure the attacker.




DATE 156–150 mya
(Late Jurassic)

SIZE 16½ ft (5 m) long

FOSSIL LOCATION Tanzania

HABITAT Forests

DIET Plants



Tuojiangosaurus

TOO-YANG-oh-SORE-us



A close relative of *Stegosaurus*, *Tuojiangosaurus* had a long, shallow snout, beaklike jaws, and spikes on its tail. It shared these features with other stegosaurs.

DATE 160–150 mya (Late Jurassic)

SIZE 23 ft (7 m) long

FOSSIL LOCATION China

HABITAT Forests

DIET Plants

Nodosaurs

A family of armored dinosaurs called the nodosaurs appeared in the Jurassic Period. Their armor was made of rows of bony deposits that formed plates and spikes on their skin. The armor helped mainly in defense but it also became important in display and combat between rivals.

Gastonia

gas-TOE-nee-ah



This low-slung and heavy nodosaur had thick, bony scutes on its back and tail, many of which were extended into bladelike spikes. If attacked, *Gastonia* could severely injure a predator with its spiked tail. The top of its skull was extra thick, suggesting that males may have had head-butting contests over territory.

DATE 125 mya (Early Cretaceous)

SIZE 13 ft (4 m) long

FOSSIL LOCATION United States

HABITAT Forests

DIET Plants



Edmontonia

ED-mon-TOE-nee-a



Edmontonia would probably scare off predators by charging and lunging at them with its spearlike shoulder spikes. It may have also used its spikes to fight with others of its kind over territory or mates.

DATE 75–65 mya (Late Cretaceous)

SIZE 23 ft (7 m) long

FOSSIL LOCATION North America

HABITAT Forests

DIET Low-growing plants

Sauropelta

SORE-oh-PELT-ah

This dinosaur's enormous neck spines helped to protect the animal against predators such as *Deinonychus* (see p. 51).

A thick shield of armor plates covered this nodosaur's back and tail, giving it the name *Sauropelta*, which means "shield lizard."


DATE 120–110 mya (Early Cretaceous)

SIZE 16½ ft (5 m) long

FOSSIL LOCATION United States

HABITAT Forests

DIET Plants



Gargyleosaurus

GAR-goil-oh-SORE-us

Unlike most other armored dinosaurs, *Gargyleosaurus* had seven conical teeth at the front of its upper jaw. These may have made it easier for the animal to tear off leaves and stems from plants. Armor plates lined the nodosaur's back, while triangular horns projected from its head and cheeks.


DATE 155–145 mya
(Late Jurassic)

SIZE 13 ft (4 m) long

FOSSIL LOCATION US

HABITAT Forests

DIET Low-growing plants



Ankylosaurs

This family of armored dinosaurs evolved in the Cretaceous Period. Unlike nodosaurs, ankylosaurs had broad, triangular heavily armored heads and lacked long spines on the sides of their body. Their tails often ended in bony clubs. The hefty clubs of some ankylosaurs may have been heavy enough to cripple predators.

Ankylosaurus

ANK-ill-oh-SORE-us

The largest of all ankylosaurs, *Ankylosaurus* was covered from head to tail with bony plates. Even its eyelids had small plates covering them. Plates at the end of the tail were fused together, forming a sledgehammerlike club, which could be swung at theropods with bone-shattering force.

DATE 70–65 mya (Late Cretaceous)

SIZE 20 ft (6 m) long

FOSSIL LOCATION North America


HABITAT Forests

DIET Plants



Bony plate on upper body

Bony club at end of tail



Minmi

MIN-mee




One of the smallest ankylosaurs, *Minmi* had extra bones along its back that may have supported its back muscles. It had small, leaf-shaped teeth and a sharp beak.

DATE

120–115 mya

(Early Cretaceous)

SIZE 10 ft (3 m) long**FOSSIL LOCATION** Australia**HABITAT** Forests and open plains**DIET** Leaves, seeds, and fruit

Euoplocephalus

YOU-owe-plo-SEFF-ah-luss



Euoplocephalus was a massive, club-tailed ankylosaur. Although a heavy animal, it had powerful legs that made it nimble on its feet. Its armor, speed, and tail club provided it with a triple defense against predators.

DATE 70–65 mya (Late Cretaceous)**SIZE** 20 ft (6 m) long**FOSSIL LOCATION** North America**HABITAT** Forests**DIET** Plants



Any predator that got past the **sledgehammer tail** of *Euoplocephalus* would be stopped by the spikes that covered its neck and back



EUOPLOCEPHALUS

In Late Cretaceous North America, *Euoplocephalus* had to fend off attacks from giant predators, such as the theropod *Gorgosaurus*. The ankylosaur's tailbones were fused together into a club, which could deliver crippling injuries to an aggressor.



FOCUS ON...

DIVERSITY

The ornithopods diversified into many different types of dinosaur.



▲ Hypsilophodonts were small, two-legged plant-eaters that could run very fast.



▲ Iguanodonts had horselike faces and ranged in size from small dinosaurs to giants.



▲ Hadrosaurs had beaks like ducks and are known as “duck-billed” dinosaurs.

Ornithopods

The ornithischians were plant-eaters with short beaks. The ornithopods made up one group of ornithischians. Some had chewing teeth that pulped plant matter. Many moved in big herds and on two legs, while some of the bigger ones usually walked on all fours.

Heterodontosaurus

HET-er-oh-DON-toe-SORE-us



Heterodontosaurus was a typical plant-eating ornithischian, except that it had three types of teeth—sharp front teeth that helped to snip off leaves, closely packed chewing teeth at the back of its mouth, and pointed, fanglike teeth, which it probably used in defense.

DATE 200–190 mya (Early Jurassic)


SIZE 3¼ ft (1 m) long

FOSSIL LOCATION South Africa

HABITAT Scrubland

DIET Plants, tubers, and possibly insects

Horny beak helped to snip off leaves



Lesothosaurus

li-SUE-too-SORE-us



This turkey-sized ornithopod was agile and could probably easily escape predators. It had big eyes on the sides of its head that gave it a good all-around view of approaching threats.

DATE 200–190 mya
(Early Jurassic)

SIZE 3¼ ft (1 m) long

FOSSIL LOCATION
South Africa

HABITAT Deserts

DIET Leaves and perhaps dead animals and insects



Dryosaurus

DRY-oh-SORE-us



Dryosaurus is the best-known member of a group of small ornithopods called dryosaurids. It could run fast and it may have flicked its tail sideways to make sharp turns for dodging obstacles or escaping predators.


DATE 155–145 mya (Late Jurassic)

SIZE 10 ft (3 m) long

FOSSIL LOCATION United States

HABITAT Forests

DIET Leaves and shoots



Leaellynasaura

lee-ELL-in-ah-SORE-ah



Leaellynasaura lived in Australia, which was much closer to the South Pole in the Cretaceous Period than it is today. This small ornithomimid faced long winters, going without sunlight for many months.

**DATE**

105 mya
(Early Cretaceous)

SIZE 6½ ft (2 m) long

FOSSIL LOCATION

Australia

HABITAT Forests

DIET Plants

Tenontosaurus

ten-NON-toe-SORE-us

Tenontosaurus had a narrow but deep skull and a rather stiff and bony tail. This animal was probably often hunted by packs of small theropods called *Deinonychus*. Teeth of this predator have been found along with *Tenontosaurus*'s bones, suggesting that the ornithomimid may have put up a fight before being brought down.

DATE 115–108 mya (Early Cretaceous)

SIZE 23 ft (7 m) long

FOSSIL LOCATION US


HABITAT Forests

DIET Plants



Hypsilophodon

HIP-sih-LOAF-oh-don



Five-fingered hand

Long, slender foot

Fossil footprints of *Hypsilophodon* suggest that it lived and moved in herds, much like deer do today. Its long legs and feet and stiff tail made it a fast-running animal, able to escape predators running only on its hind legs and balancing with its tail.

DATE 125–120 mya (Early Cretaceous)

SIZE 6½ ft (2 m) long

FOSSIL LOCATIONS England and Spain

HABITAT Forests

DIET Plants




The long, pointed claws on this dinosaur's feet gave it a dangerous kick.

Muttaborrasaurus

MOO-tah-BUH-ruh-SORE-us

This ornithopod may have used its huge, hollow nose to create honking sounds or to warm the cool air that the animal breathed in. *Muttaborrasaurus* had strong skull bones beneath its eye sockets, which allowed it to bite and chew tough plants.

Enlarged, bony growth on snout



DATE 100–98 mya (Early Cretaceous)

SIZE 26 ft (8 m) long

FOSSIL LOCATION Australia

HABITAT Forests

DIET Plants

Lesothosaurus was
built like a gazelle
and could easily escape the
bulky predators of the time




**LESOTHO SAURUS**

Lesothosaurus lived during the Early Jurassic and was among the earliest ornithomids. It was agile and is seen here evading the crocodilelike predator *Sphenosuchus*. *Lesothosaurus*'s small teeth were shaped like arrowheads, and it probably grazed on low-growing plants.

Rhabdodon

RAB-doe-don



Although it was discovered in 1869, it is still unclear whether *Rhabdodon* belonged to the hypsilophodont family or the iguanodonts.

This broad-bodied ornithopod had stout jaw bones and rounded teeth.

DATE 75 mya (Late Cretaceous)

SIZE 12 ft (3.7 m) long

FOSSIL LOCATIONS Austria, France, Romania, and Spain

HABITAT Forests

DIET Plants

Camptosaurus

CAMP-toe-SORE-us



Camptosaurus belonged to the group of ornithopods known as iguanodonts. It was a heavily built animal with a long, horselike face tipped by a beak. Its hands had a thumb spike and the middle fingers supported the weight of the animal's body when on all fours.


DATE 155–145 mya (Late Jurassic)

SIZE 16½ ft (5 m) long

FOSSIL LOCATION United States

HABITAT Forests

DIET Low-growing herbs and shrubs



Iguanodon

ig-GWAH-no-don

Iguanodon's long jaws had leaf-shaped teeth similar to those of a modern-day iguana. It spent most of its time on four legs, feeding on the ground, but could rear up on its hind legs to reach for food on trees. Strong middle toes on its feet supported its weight. In 1825, it became the second prehistoric animal to be identified as a dinosaur, after *Megalosaurus*.


DATE 135–125 mya (Early Cretaceous)

SIZE 29½–39 ft (9–12 m) long

FOSSIL LOCATIONS Belgium, Germany, France, Spain, and England

HABITAT Forests

DIET Plants



Maiasaura

MY-a-SORE-a

A fossil site in Montana has remains of numerous bowl-shaped dinosaur nests close to each other. Scientists believe that this was a nesting colony, much like the colonies of modern seabirds, where parents raised their young. They named the nest builders *Maiasaura*, or “good mother lizard.”



DATE 80–74 mya (Late Cretaceous)

SIZE 29½ ft (9 m) long

FOSSIL LOCATION United States

HABITAT Coastal plains

DIET Leaves

Hadrosaurus

HAD-roh-SORE-us

This ornithomimid used its toothless, beak for tearing twigs and leaves from plants. It had hundreds of blunt teeth at the back of its mouth that ground its food to a pulp.


DATE 80–74 mya (Late Cretaceous)

SIZE 29½ ft (9 m) long

FOSSIL LOCATION North America

HABITAT Forests

DIET Leaves and twigs



Corythosaurus

ko-RITH-oh-SORE-us

The crest of this hadrosaur, or duck-billed dinosaur, looks similar to the helmets worn by the soldiers of Corinth in ancient Greece. This inspired its name. Its crest probably functioned like a trombone, helping it make loud, booming calls to other members of its herd.

DATE 76–74 mya (Late Cretaceous)

SIZE 29½ ft (9 m) long

FOSSIL LOCATION Canada

HABITAT Forests and swampy areas

DIET Pine needles and seeds

Lambeosaurus

LAMB-ee-oh-SORE-us

This skeleton is mounted to show how the animal might have stooped to drink from a pond or river

Hatchet-shaped crest

Long pubis (hip bone)

The hollow crest on this hadrosaur's head changed shape as it grew into an adult. The crest probably helped it to recognize others of its kind. Males may have used it to impress females.

DATE 76–74 mya (Late Cretaceous)

SIZE 29½–49 ft (9–15 m) long

FOSSIL LOCATION Canada

HABITAT Forests

DIET Low-growing leaves, fruits, and seeds



Edmontosaurus

ed-MONT-oh-SORE-us

This duck-billed dinosaur had a broad beak, which it used to crop leaves. Then it chewed the food to a pulp with more than 1,000 tiny cheek teeth. Like other hadrosaurs, its hind legs were longer than its forelegs, but it spent most of the time on all fours.





DATE 75–65 mya (Late Cretaceous)

SIZE 43 ft (13 m) long

FOSSIL LOCATIONS US and Canada

HABITAT Swamps

DIET Plants



Ducklike beak

Brachylophosaurus

BRACK-ee-LOAF-oh-SORE-us



Brachylophosaurus had a rectangular skull with a flat, paddle-shaped bony crest on its head. The crest was much wider and more heavily built in males than in females.



DATE 76.5 mya (Late Cretaceous)

SIZE 29½ ft (9 m) long

FOSSIL LOCATION North America

HABITAT Forests

DIET Ferns, flowering plants, and conifers

Parasaurolophus

PA-ra-SORE-oh-LOAF-us



The most striking feature of this dinosaur was the tubelike crest on its head. The cavity within was linked to the animal's nostrils, and *Parasaurolophus* may have used it to make loud, trumpetlike calls to communicate with its herd members.


DATE 76–74 mya (Late Cretaceous)

SIZE 29½ ft (9 m) long

FOSSIL LOCATION North America

HABITAT Forests

DIET Pine needles and seeds



Pachycephalosaurs

One of the last groups of dinosaur to evolve was the pachycephalosaurs, or “thick-headed lizards.” These dinosaurs get their name from the thick domes on the tops of their skulls. They had many kinds of small, ridged teeth that helped to shred leaves and other vegetation.

Stegoceras

steg-OSS-er-us

This plant-eater had coarsely serrated teeth, probably used for tearing and chewing leaves. Its skull not only had a domed roof, but also a bony shelf at the back, dotted with knobs and spikes.

DATE 77.5–74 mya (Late Cretaceous)

SIZE 6½ ft (2 m) long

FOSSIL LOCATION Canada

HABITAT Forests


DIET Leaves and fruits



Bony shelf on domed skull



Stegoceras's skull was shaped like a motorcycle helmet and was about 3½ in (9 cm) thick.



Pachycephalosaur

PACK-ee-sef-ah-low-SORE-us



The thick skull of this pachycephalosaur was ringed with a crown of bony spikes. The spikes also lined its cheeks and muzzle. The function of this headgear is unknown—rivals may have butted heads or it may have been used to impress mates. *Pachycephalosaur* had leaflike teeth at the side of its mouth, nipping teeth at front, and conical ones in its lower jaw.

DATE 65 mya (Late Cretaceous)

SIZE 16½ ft (5 m) long

FOSSIL LOCATION North America

HABITAT Forests

DIET Plants, soft fruits, and seeds





FOCUS ON...

HORNS

Horns evolved from small knobs to formidable weapons.



▲ Early ceratopsians, such as *Psittacosaurus*, had small, bony, hornlike growths on their cheeks.



▲ The main horn of *Centrosaurus* grew over its nose and it used it for defense.



▲ Horns above the eyes not only made *Triceratops* look fierce, but also served as weapons in combat.

Ceratopsians


Although ceratopsians were plant-eaters, their long horns and frills gave them a formidable appearance. Living in herds, they grazed the forests and plains of North America and Asia.

Psittacosaurus

SIT-ack-oh-SORE-us



One of the earliest members of the ceratopsians, *Psittacosaurus* had strong hind legs, which suggest that it could run quickly on only two legs.



DATE 120–100 mya
(Early Cretaceous)

SIZE 6½ ft (2 m) long

FOSSIL LOCATIONS

China and Mongolia


HABITAT Deserts and scrublands

DIET Plants

The dinosaur is asleep in this reconstruction

Protoceratops

PRO-toe-SERRA-tops



A small ceratopsian, *Protoceratops* had broad feet and wide, spadelike claws that allowed it to dig burrows for shelter from the desert Sun.

DATE 74–65 mya (Late Cretaceous)

SIZE 6 ft (1.8 m) long

FOSSIL LOCATION Mongolia

HABITAT Deserts

DIET Desert plants

Triceratops

try-SERRA-tops



As heavy as a 10-ton truck, this dinosaur was built like a modern rhinoceros. *Tyrannosaurus* bite marks found on *Triceratops* skulls suggest that there were fierce battles between these two species of dinosaur.

DATE 70–65 mya (Late Cretaceous)

SIZE 29½ ft (9 m) long

FOSSIL LOCATION
North America

HABITAT Forests

DIET Forest plants



Pentaceratops

PEN-ta-SERRA-tops



The most remarkable feature of this dinosaur was its huge head. One fossil skull, built from broken fragments, is more than 10 ft (3 m) long, making it the longest skull of any land animal in history. *Pentaceratops* had five horns on its face, one on the snout, one curved horn on each eyebrow, and a small horn on each cheek.

DATE 76–74 mya (Late Cretaceous)

SIZE 21½ ft (6.5 m) long

FOSSIL LOCATION United States

HABITAT Forests

DIET Plants

Eye spots may have enhanced the animal's neck frill display



Chasmosaurus

KAZ-mo-SORE-us

An enormous neck frill, which reached over the shoulders, was the most distinctive feature of this dinosaur. Holes present in the frill were covered with brightly colored skin and helped attract mates. *Chasmosaurus* could also tilt its frill upright and startle enemies and predators.

DATE 74–65 mya (Late Cretaceous)

SIZE 16½ ft (5 m) long

FOSSIL LOCATION North America

HABITAT Forests

DIET Palms and cycads



Einiosaurus

EYE-nee-oh-SORE-us

The front horn of this dinosaur was very different from that of other ceratopsians. In a young *Einiosaurus*, the horn was straight, but as the animal grew, it gradually curved forward. These dinosaurs lived in herds, moving from place to place to find fresh grazing.

DATE 74–65 mya (Late Cretaceous)

SIZE 20 ft (6 m) long

FOSSIL LOCATION US

HABITAT Forests

DIET Plants





Styracosaurus

sty-RACK-oh-SORE-us



Six spikes decorated the frill of this spectacular reptile. The spikes on a male served as a decoration, which attracted females, and the longer the spikes, the more attractive the male looked. The teeth of a *Styracosaurus* grew constantly, replacing worn ones, as the dinosaur munched through the Cretaceous forests.

DATE 74–65 mya (Late Cretaceous)

SIZE 17 ft (5.2 m) long

FOSSIL LOCATION North America

HABITAT Open woodlands

DIET Ferns and cycads





Dinosaur neighbors

Dinosaurs were not the only land animals that lived in the Mesozoic Era. They shared the planet with many other creatures. These included other archosaurs, and the rhynchosaurs, cynodonts, and early mammals. These dinosaur neighbors ranged in size from the small, shrewlike mammal *Eomaia* to the dinosaur-sized, bipedal archosaur *Postosuchus* (left).



EFFIGIA

Many archosaurs, such as *Effigia*, were similar to dinosaurs, but are more closely related to modern crocodiles and alligators.

Rhynchosaurs

These barrel-shaped plant-eaters outnumbered dinosaurs during the Triassic. They had a beak at the front of their mouth and several rows of teeth on the mouth's roof. They used tusks to shear plants and then ground and crushed the plant matter before swallowing it.

Rhynchosaurus

RIN-cho-SORE-us

This reptile had a typical rhynchosaur beak and a deep, broad lower jaw. The skeletons found show that it was well-adapted for fast movement across the ground and had semierect hind limbs. *Rhynchosaurus* used its hind feet to dig out roots and tubers from the soil.

DATE 245–240 mya (Middle Triassic)

SIZE 1½–3¼ ft (0.5–1 m) long

FOSSIL LOCATION England

HABITAT Semiarid plains


DIET Ferns and tubers



More than 35 skeletons of *Hyperodapedon* have been found in rocks 230 million years old in Elgin, Scotland.

Tail dragged on the ground





Razor-sharp beak

Hyperodapedon

HIGH-per-oh-DAP-eh-don



Like other members of its group, *Hyperodapedon* was a heavy-bodied, four-legged plant-eater.

It had a pig-shaped body and a relatively long tail and its head was large and deep. Its beak was curved and it had two short tusks.

DATE 231–216 mya (Late Triassic)

SIZE 4–5 ft (1.2–1.5 m) long

FOSSIL LOCATIONS Scotland, Argentina, Brazil, and India

HABITAT Forests

DIET Seed ferns and other plants

Archosaurs

The first archosaurs evolved around 255 million years ago. From these, an assortment of different animals evolved, including crocodylians, pterosaurs, and dinosaurs.



FOCUS ON...

DIVERSITY

The crocodylians and their relatives were called the crurotarsans.

Stagonolepis

stag-o-NO-lep-iss



Shovel-like snout

Stagonolepis was one of a group of heavily armored archosaurs called the aetosaurs. Bony armor plates ran the length of its body. It had a short, deep skull that ended in a shovel-like snout, which probably helped it to dig out juicy roots like a pig does today.

DATE 235–223 mya (Late Triassic)

SIZE 10 ft (3 m) long

FOSSIL LOCATIONS Scotland, Poland, and South America

HABITAT Forests

DIET Horsetails, ferns, and cycads



▲ Many crurotarsans, such as the crocodilelike *Deinosuchus*, had sprawling limbs.



▲ Others, such as the dinosaurlike *Postosuchus*, could walk on their upright hind limbs.



▲ *Effigia* was even more dinosaurlike, closely resembling the ostrichlike theropods.

Desmosuchus

des-MAT-o-SOOK-us

Desmosuchus was an aetosaur that resembled a short-snouted crocodile. It had rows of rectangular bony plates along its back and tail. The underside of part of its belly was also covered with these plates. The spikes on its shoulders could reach a length of 18 in (45 cm).

DATE 230 mya
(Late Triassic)

SIZE 16½ ft (5 m) long

FOSSIL LOCATION
United States


HABITAT Forests

DIET Plants




Bony plates
on back

Toothless
snout



Lagosuchus

LAG-o-SOOK-us



Lagosuchus was an agile archosaur with long, thin hind legs and long feet. These enabled it to chase its prey at great speeds or escape predators quickly.

DATE 230 mya (Late Triassic)

SIZE 12 in (30 cm)

FOSSIL LOCATION Argentina

HABITAT Forests

DIET Small animals

Parasuchus

para-SOOK-us



Parasuchus was a phytosaur—a type of long-snouted archosaur that spent a lot of time in water and resembled today's crocodiles. It hunted for prey near the water's edge. Its eyes faced sideways.



DATE 225 mya (Late Triassic)

SIZE 6½ ft (2 m) long

FOSSIL LOCATION India

HABITAT Rivers and swamps

DIET Fish and small reptiles

Effigia

eff-IDGE-ee-ah



Effigia was a raiusuchian—a type of archosaur with upright legs. Raiusuchians evolved in the Triassic Period. *Effigia* was an omnivore and had a toothless beak like many dinosaurs. It may have used it to crack open seeds and eggs, shear vegetation, or feed on small animals. *Effigia* means “ghost” in Greek—the creature was named after the Ghost Ranch Quarry in New Mexico, where its fossils were found in 1947.

DATE 210 mya (Late Triassic)

SIZE 6½–10 ft (2–3 m) long

FOSSIL LOCATION United States

HABITAT Forests

DIET Plants, seeds, and animals

Postosuchus

POST-oh-SOOK-us





Postosuchus was a large rauisuchian and one of the largest predators of its time. Its large skull had curved, daggerlike teeth that looked like those of the large theropods. It lived alongside the first dinosaurs and probably preyed on them.

DATE 230–200 mya (Late Triassic to Early Jurassic)

SIZE 14¾ ft (4.5 m) long

FOSSIL LOCATION United States

HABITAT Forests


DIET Small reptiles



Ornithosuchus

or-nith-oh-SOOK-us

Although it mostly moved around on four limbs, this early archosaur could probably also walk and run on two legs. It used its sharp teeth to slice through the flesh of prey.



DATE 230 mya (Late Triassic)

SIZE 13 ft (4 m) long


FOSSIL LOCATION Scotland

HABITAT Swamps of western Europe

DIET Small animals

Dakosaurus

DACK-oh-SORE-us



This distant relative of crocodiles was part of a group of fierce marine predators called the metriorhynchids. Its deep skull resembled that of *Tyrannosaurus*. The skull was lined with sharp teeth that helped *Dakosaurus* to slice through the flesh of other reptiles and crunch the shells of ammonites.

DATE 165–140 mya (Late Jurassic)

SIZE 13–16½ ft (4–5 m) long

FOSSIL LOCATIONS Western Europe, Mexico, and Argentina

HABITAT Shallow seas

DIET Fish, ammonites, and marine reptiles

Terrestriusuchus

teh-REST-rih-SOO-kuss



This tiny carnivore had pencil-thin limb bones and a lightweight skull. It walked with its body raised off the ground. Like modern crocodiles, it had bony plates on its skin.



DATE 215–200 mya (Late Triassic)

SIZE 2½–3¼ ft (0.75–1 m) long

FOSSIL LOCATIONS British Isles, W. Europe

HABITAT Dry uplands and forests

DIET Insects and small animals

Sphenosuchus

SFEN-oh-SOO-kuss



This slender-legged creature could probably run fast when fleeing predators or chasing prey. Parts of its skull had air-filled spaces.

DATE 200 mya (Early Jurassic)

SIZE 3¼–5 ft (1–1.5 m) long

FOSSIL LOCATION South Africa

HABITAT Banks of rivers and lakes in humid lowlands

DIET Small land animals

Deinosuchus

DYE-no-SOO-kuss

*Platelike scales
made of bone*



Deinosuchus was nearly five times bigger than any alligator found today. It may have lurked patiently at the water's edge, waiting to pounce on passing fish, marine reptiles, or even dinosaurs as large as itself. Much like modern alligators do, it dragged its victims under water and drowned them.

DATE 70–65 mya (Late Cretaceous)

SIZE 33 ft (10 m) long

FOSSIL LOCATIONS United States and Mexico

HABITAT Swamps

DIET Fish and medium to large dinosaurs

Simosuchus

SIGH-moe-SOO-kuss

The name of this creature means “pug-nosed crocodile.” Its short skull and blunt face were unusual for a crocodylian. Its teeth suggest that it may have fed mainly on plants. Its hind limbs were semierect and it probably did not run.


DATE 70 mya (Late Cretaceous)

SIZE 4 ft (1.2 m) long

FOSSIL LOCATION Madagascar

HABITAT Forests


DIET Plants and maybe some insects



**These animals looked like dinosaurs
and moved like dinosaurs, but they are**

imposters —

**members of an earlier group
of reptiles called rauisuchians**



**EFFIGIA**

Effigia looked just like a type of theropod called an ostrich dinosaur. Actually, it was a kind of archosaur called a raiusuchian and lived 80 million years earlier. Like an ostrich dinosaur, it ran on its hind legs and balanced with its long tail.

Cynodonts and dicynodonts

The cynodonts formed a group of mammal-like reptiles that included the ancestors of modern mammals. Their bodies may have been covered in hair and they walked on upright legs. The cynodonts lived alongside the dicynodonts, which formed another group of mammal-like reptiles with two tusks and a blunt beak.



FOCUS ON... TEETH

Both cynodonts and dicynodonts had distinctive teeth.



▲ Cynodont means “dog teeth” — these creatures evolved mammal-like teeth.



▲ Dicynodonts were distinguished by a pair of tusklike canines in the front of the mouth.

Placerias

plah-SEE-ree-ass



Dicynodonts were plant-eaters. *Placerias* was the largest dicynodont in its environment. It looked like a hippopotamus and weighed about 1,300 lb (600 kg). It used its horny beak to shear plants.

DATE 220–215 mya (Late Triassic)

SIZE 6½–11½ ft (2–3.5 m) long

FOSSIL LOCATION United States

HABITAT Flood plains

DIET Plants




Lystrosaurus

Lis-trow-SORE-us



Many species of land-living animals died out at the end of the Permian Period. This dicynodont was one of the few animals to survive. *Lystrosaurus* had a piglike, barrel-chested body. Like all dicynodonts, it probably used its canines for display or in defense.



DATE 255–230 mya
(Late Permian–Late Triassic)

SIZE 3¼ ft (1 m) long

FOSSIL LOCATIONS Africa, Russia, India, China, Mongolia, and Antarctica

HABITAT Dry flood plains

DIET Plants

Cynognathus

SIGH-nog-NA-thus



The name of this wolf-sized cynodont means “dog-jaw,” and it had a large doglike canine on each side of its jaw. It also had bladelike incisors that helped slice flesh. Cynodonts used molar teeth for chewing food, much like their mammal descendants.

DATE 247–237 mya (Early–Middle Triassic)

SIZE 3¼ ft (1 m) long

FOSSIL LOCATIONS South Africa, Antarctica, and Argentina

HABITAT Forests

DIET Meat



Thrinaxodon

thrin-AX-oh-don



This catlike predator was the most common cynodont in the early Triassic. It held its limbs almost under its body, like modern mammals do, and it may have had a fur covering.

DATE 248–245 mya (Early Triassic)

SIZE 12 in (30 cm) long


FOSSIL LOCATIONS

South Africa and Antarctica

HABITAT

Burrows in forests and riverbanks

DIET Insects and reptiles



Early mammals

The first mammals evolved in the Triassic Period from cynodont ancestors. These shrewlike early mammals were probably furry and warm-blooded like today's mammals, and many of them may have had a good sense of smell. The early mammals lived alongside the dinosaurs, and the largest of them fed on baby dinosaurs.

Morganucodon

MORE-gan-YOU-koh-don



This tiny, shrewlike animal was one of the first true mammals. *Morganucodon* had several features typical of its reptilian ancestors, including a double jaw joint. It probably laid eggs as reptiles do and actively hunted at night. It was discovered in 1949 in Wales.

DATE 210–180 mya (Late Triassic–Early Jurassic)

SIZE 3½ in (9 cm) long

FOSSIL LOCATIONS Wales, China, and US

HABITAT Forests

DIET Insects

Sharp teeth



Nemegtbaatar

nem-EGT-bat-or



Wide snout

Nemegtbaatar looked like a vole because of its short, deep skull. It had a wide snout and its front teeth juttied out, giving it a buck-toothed face. It may have been a plant-eater.

DATE 65 mya (Late Cretaceous)

SIZE 4 in (10 cm) long

FOSSIL LOCATION Mongolia

HABITAT Forests

DIET Possibly plants

Megazostrodon

MEG-ah-ZO-stroh-don



This animal had a slender body, with a long snout and tail. It probably burrowed and ran like today's rats and shrews. Each of its cheek teeth had short, triangular points, possibly used for cutting up insects.



DATE 190 mya (Early Jurassic)

SIZE 4 in (10 cm) long

FOSSIL LOCATION South Africa

HABITAT Forests

DIET Insects

Eomaia

EE-oh-MY-ah



Rat-sized *Eomaia*'s name stands for "dawn mother." It was one of the first placental mammals, the large group of modern mammals with a placenta—a structure inside the mother's body that nourishes the developing young.

DATE 125 mya (Early Cretaceous)

SIZE 8 in (20 cm) long

FOSSIL LOCATION China

HABITAT Forests

DIET Insects and other small animals



Sinoconodon

SIGH-no-CON-oh-don



Sinoconodon was the size of a squirrel. It had a strong jaw joint and chin and may have had a powerful bite. Its ear bones were like those of a mammal, but its teeth were replaced throughout life, as in a reptile.

DATE 200 mya
(Early Jurassic)

SIZE 12 in (30 cm) long

FOSSIL LOCATION China

HABITAT Forests

DIET Omnivorous





Sea reptiles

While dinosaurs were the dominant life forms on land, the oceans were ruled by gigantic predatory reptiles, which included the plesiosaurs, nothosaurs, and mosasaurs (left). These reptiles used their flipper-shaped limbs to surge rapidly through the ocean, often chasing prey. Many of them had pointed teeth that held on to slippery fish.



TURTLES

Aside from monstrous predators, other marine reptiles included the placodonts and turtles, like *Protostega*, which developed thick defensive plates on its back.

Placodonts and turtles

During the Middle Triassic Period, shallow coastal seas covering Europe teemed with predatory reptiles called placodonts. They had a large, barrel-chested body, webbed limbs that worked as paddles, and a long, deep tail. The earliest turtles also lived around the same time. They had a short skull, a tiny tail, and a protective shell.

Placodus

plak-OH-dus



This reptile was an expert swimmer, despite its bulky body. Unusually for a reptile, it had protruding front teeth that probably speared fish. Peglike teeth on the roof of its mouth would have crushed the hard shells of mollusks.


DATE 245–235 mya (Middle Triassic)

SIZE 6½–10 ft (2–3 m) long

FOSSIL LOCATION Germany

HABITAT Shallow seas near reefs

DIET Mussels, fish, and other invertebrates



Kayentachelys

ka-YEN-ta-KEE-lis



Kayentachelys was among the first turtles to evolve the fully formed, boxlike protective shell seen in modern species. It had a sharp beak like its modern relatives and like most of today's turtles, it could pull in its head and limbs to hide within the protective shell, if needed.

DATE 196–183 mya (Early Jurassic)

SIZE 23½ in (60 cm) long


FOSSIL LOCATION United States

HABITAT Near streams in arid regions

DIET Plants and animals

Odontochelys

oh-DON-toh-KEE-lis

Odontochelys is the most ancient and primitive turtle discovered so far.

Odontochelys differed from modern turtles in two major ways. Today's turtles have a toothless beak, while this ancestral creature had jaws lined with teeth. This inspired its name, which means "toothed turtle." It also had a shell that covered only its belly, unlike those of modern turtles, which protect both the belly and the back.

DATE 220 mya (Late Triassic)

SIZE 16 in (40 cm) long

FOSSIL LOCATION China

HABITAT Shallow coastal seas

DIET Fish, ammonites, and plants

Nothosaurs

The nothosaurs patrolled the shallow coastal seas of the Triassic at about the same time as the first dinosaurs appeared on land. These fish-hunters had four legs with webbed feet and probably bled on beaches and coastal rocks, much like seals do today.

Pachypleurosaurus

PACK-ee-ploo-roe-SORE-us



Pachypleurosaurus was a small animal with a snakelike, streamlined body and a long tail. It swam like an otter—by moving its body in a wavelike pattern—and steered with the help of paddlelike hind limbs.

DATE 225 mya
(Late Triassic)

SIZE 12–16 in
(30–40 cm) long

FOSSIL LOCATIONS
Italy and Switzerland

HABITAT
Shallow oceans

DIET Small fish



Lariosaurus

LA-ree-oh-SORE-us



A small nothosaur, *Lariosaurus* lived mainly in water, but often came out onto land. Unlike most reptiles, this nothosaur probably gave birth to live young.

DATE 234–227 mya (Late Triassic)

SIZE 20–28 in (50–70 cm) long

FOSSIL LOCATION Italy

HABITAT Shallow oceans

DIET Small fish and shrimp



Nothosaurus

NO-tho-SORE-us



Like other nothosaurs, *Nothosaurus* probably swam by waving its long body and tail to move through water. Its long, pointed teeth were good at gripping slippery prey. Much like crocodiles, it could throw its head to one side to seize passing fish.


DATE 240–210 mya (Middle–Late Triassic)

SIZE 4–13 ft (1.2–4 m) long

FOSSIL LOCATIONS Europe, Africa, Russia, and China

HABITAT Shallow oceans

DIET Fish and shrimp



Ichthyosaurs

The oceans of the Mesozoic Era were home to many predatory reptiles, including this group of dolphinlike beasts. Streamlined hunters, the ichthyosaurs swam using their sharklike fins and tails and fed on squid, ammonites, fish, and marine reptiles. They had large eyes and gave birth to young in water.

Shonisaurus
SHON-ee-SORE-us



Shonisaurus had a long, toothless snout. When catching prey, strong muscles in the mouth pulled its tongue back rapidly, helping to suck in its victims. It was the largest marine reptile ever to live on the Earth.

DATE 225–208 mya (Late Triassic)

SIZE Up to 69 ft (21 m) long


FOSSIL LOCATION North America

HABITAT Open oceans

DIET Fish and squid

Ichthyosaurus

ICK-three-oh-SORE-us



A small predator, *Ichthyosaurus* could catch fast-moving, slippery prey—its snout was long and armed with banks of sharp, needlelike teeth. Like all ichthyosaurs, it hunted mainly using eyesight. Its large eyes were protected by bony shields.

DATE 190 mya (Early Jurassic)

SIZE 6½ ft (2 m) long

FOSSIL LOCATIONS British Isles, Belgium, and Germany

HABITAT Open oceans

DIET Fish and squid

Mixosaurus

MIX-oh-SORE-us



Mixosaurus was an early ichthyosaur and like others of its kind, it swam by beating its tail from side to side, probably using bursts of speed to chase after or surge through shoals of fish.

DATE 230 mya (Late Triassic)

SIZE Up to 3¼ ft (1 m) long

FOSSIL LOCATIONS North America, Europe, Asia

HABITAT Open oceans

DIET Fish

Temnodontosaurus

tem-NOH-don-toh-SORE-us



This large ichthyosaur could dive to great depths in the seas when hunting for prey. Its eyes had a diameter of 8 in (20 cm), making them larger than those of most other vertebrates.

DATE 198–185 mya (Early Jurassic)

SIZE 39 ft (12 m) long

FOSSIL LOCATIONS England and Germany

HABITAT Shallow seas

DIET Fish
and squid




Plesiosaurs

Fully adapted to an aquatic life, these gigantic carnivorous reptiles dominated the oceans in the Jurassic and Cretaceous periods, swimming in the water with four long flippers. Many of them had long, snakelike necks and small heads.



Cryptoclidus

KRIP-toe-KLIDE-us



This plesiosaur had a flat head and a light skull. It had hundreds of teeth, which interlocked with each other and trapped fish and other small marine animals. Like other plesiosaurs, it probably “flew” through water, moving its flippers like wings. Some scientists think that it may have come ashore to lay eggs.

DATE 165–150 mya (Middle to Late Jurassic)

SIZE 26 ft (8 m) long

FOSSIL LOCATIONS England, France, Russia, and South America

HABITAT Shallow oceans

DIET Fish and squid

Elasmosaurus

el-LAZZ-moe-SORE-us

A fish-eater, *Elasmosaurus* probably swam over the seabed, reaching down to grab prey. Its neck was supported by 72 vertebrae (spine bones)—more than that in any other animal that has ever lived.


DATE 99–65 mya (Late Cretaceous)

SIZE 46 ft (14 m) long

FOSSIL LOCATION United States

HABITAT Open oceans

DIET Fish, squid, and ammonites



Plesiosaurus

PLEE-see-oh-SORE-us

Plesiosaurus had a wide, turtlelike body. It may have hunted by swimming among shoals of fish, swinging its long neck from side to side to catch prey. It had wide, U-shaped jaws and sharp, conical teeth that gripped prey.


DATE 200 mya (Early Jurassic)

SIZE 10–16½ ft (3–5 m) long

FOSSIL LOCATIONS British Isles and Germany

HABITAT Open oceans

DIET Fish and ammonites



Pliosaurus

Plesiosaurs with short necks and large heads, the pliosaur were some of the most formidable predators ever to swim in the Earth's oceans. With muscular necks, huge jaws, and crocodilelike teeth, these sea monsters attacked and ate any creature they swam into. Their main foes were gigantic sharks and others of their own kind.

Liopleurodon

LIE-oh-PLOOR-oh-don



Liopleurodon's jaws were very powerful and it may have had a stronger bite than *Tyrannosaurus*. Scientists think that this creature had a keen sense of smell that enabled it to hunt in deep waters where prey was difficult to spot. Its long, paddlelike flippers suggest that it swam at high speeds in short spurts.

DATE 165–150 mya (Middle to Late Jurassic)

SIZE 16½–23 ft (5–7 m) long

FOSSIL LOCATIONS British Isles, Germany, France, and Russia

HABITAT Open oceans

DIET Large squid and ichthyosaurs

Rhomaleosaurus

ROME-alley-oh-SORE-us

This pliosaur had a good sense of smell and could pick up the scent of prey from far away. It also had sharp eyesight, allowing it to hunt at close range. Once it had seized prey, it probably twisted around in the water to rip its victim apart, much like crocodiles do.

DATE 200–195 mya (Early Jurassic)

SIZE 16½–23 ft (5–7 m) long

FOSSIL LOCATIONS England and Germany

HABITAT Coastal waters

DIET Fish, squid, and ocean reptiles



Kronosaurus

crow-no-SORE-us

Like a modern crocodile, *Kronosaurus* could open its jaws wide to seize prey. The head of this monstrous predator was about 10 ft (3 m) long and larger than a man.

Its skull was nearly twice the size of a *Tyrannosaurus* skull.


DATE 65 mya (Late Cretaceous)

SIZE 33 ft (10 m) long

FOSSIL LOCATIONS Australia and Colombia

HABITAT Open oceans

DIET Marine reptiles, fish, and ammonites



Rhomaleosaurus
rushed in to
cripple
its prey
with a series of massive
bites, rather like a
great white shark



**RHOMAELOS AURUS**


Rhomaelosaurus was a plesiosaur that preyed on fish, ichthyosaurs, and smaller plesiosaur. Like all plesiosaurs, it could smell prey from far away by channeling water to sense organs above the roof of its mouth. These detected scents, then the water drained out through its nostrils.

Mosasaurus

Colossal lizards called mosasaurs were major predators in Late Cretaceous seas. They evolved from small land-dwelling lizards that took to the water. Fully adapted to marine life, they had paddlelike limbs and swam like crocodiles.

Mosasaurus

MOZE-ah-SORE-us



This crocodilelike hunter swam by moving its long body in slow waves and probably hunted slow-moving prey in the surface waters of the oceans. It could even overpower plesiosaurs.


DATE 70–65 mya (Late Cretaceous)

SIZE 49–57½ ft (15–17.6 m) long

FOSSIL LOCATIONS US, Belgium, Japan, Netherlands, New Zealand, Morocco, and Turkey

HABITAT Shallow coastal oceans

DIET Fish, squid, plesiosaurs, and shellfish



Plioplatecarpus

PLEE-o-PLAH-teh-CAR-pus



This mid-sized predator preferred warm and shallow oceans. The shape of its teeth and skull suggests that it probably hunted small prey. Its long skull was equipped with thick, conical teeth. It had larger eyes than other mosasaurs.


DATE 83.5 mya (Late Cretaceous)

SIZE 16½–20 ft (5–6 m) long

FOSSIL LOCATIONS Europe, Canada, and United States

HABITAT Shallow oceans

DIET Fish





Flying reptiles

The Mesozoic Era saw the evolution and extinction of some extraordinary flying reptiles. These were the pterosaurs, which first took to the skies in the Triassic and by the end of the Cretaceous had reached colossal proportions. Pterosaurs included the agile *Pterodactylus*, which snatched fish out of the oceans as well as the gigantic *Quetzalcoatlus* (left), which stalked through Cretaceous forests, preying on dinosaurs.



PTERODACTYLUS

A Jurassic pterosaur, *Pterodactylus* lived near coasts, hunting during the day and sleeping at night.

Pterosaurs

The reptiles that took to the air in the Triassic were a type of archosaur called pterosaurs and some were the largest creatures ever to fly. They had batlike wings made of skin and their bodies were covered in fur.



FOCUS ON... TAILS

The aerodynamics of pterosaurs improved during their evolution.

Rhamphorhynchus

ram-foe-RINK-us

This pterosaur had a long, bony tail. The tail had a diamond-shaped flap of skin at the end that may have worked like a rudder, helping this reptile steer.


DATE 150 mya (Late Jurassic)

SIZE 1–6 ft (0.3–1.8 m) wingspan

FOSSIL LOCATIONS Europe and Africa

HABITAT Coasts and riverbanks

DIET Fish





▲ Typical Triassic pterosaurs, such as *Eudimorphodon*, had long tails and short legs and wings. They belonged to a group called the rhamphorhynchoids.



▲ A new group called the pterodactyloids evolved in the Jurassic. Their shorter tails and longer wings made them more agile in the air.



Dimorphodon

die-MORE-foe-don



This creature's huge head was almost one-third of its body length. Unusually for a pterosaur, it had two types of teeth. Longer teeth in the front helped snap up prey, while the ones at the back were used to grind food.

DATE 200–180 mya (Early Jurassic)

SIZE 4¾ ft (1.45 m) wingspan

FOSSIL LOCATION British Isles


HABITAT Coastal forests

DIET Fish and small, lizardlike reptiles

Peteinosaurus

pet-INE-oh-SORE-us

This reptile's name means "winged lizard." The wings of this early pterosaur were far smaller than those of the Cretaceous pterosaurs, such as *Pteranodon*.



DATE 228–215 mya (Late Triassic)

SIZE 23½ in (60 cm) wingspan

FOSSIL LOCATION Italy

HABITAT Swamps and river valleys

DIET Flying insects

Anurognathus

an-YOOR-og-NATH-us

This tiny pterosaur may have fed on damselflies and lacewings. It probably landed on the backs of the sauropods before attacking the insects flying near them.




DATE 150–145 mya (Late Jurassic)

SIZE 20 in (50.8 cm) wingspan

FOSSIL LOCATION Germany

HABITAT Forests

DIET Flying insects



The small crest on *Pterodactylus*'s head was made from bone and thick skin and was probably used for display.

Pterodactylus

TEH-roe-DACK-till-us

Many complete fossils of *Pterodactylus* have been found, making it the best-known of all pterosaurs. It had a much smaller tail and longer wing bones than earlier pterosaurs, making it far more agile in flight.

DATE 150–144 mya (Late Jurassic)

SIZE 3¼ ft (1 m) wingspan

FOSSIL LOCATION Germany

HABITAT Coastal areas

DIET Fish, insects, and perhaps carrion



Quetzalcoatlus

KWETS-ul-coe-AT-luss



Named after the Aztec god Quetzalcoatl, this monstrous pterosaur stood as tall as a giraffe. Besides soaring in the air like a condor, *Quetzalcoatlus* could also hunt for prey like a giant stork, walking on land and picking up small dinosaurs with its beak.

DATE 70–65 mya (Late Cretaceous)

SIZE 33–36 ft (10–11 m) wingspan

FOSSIL LOCATION United States

HABITAT Open plains and forests

DIET Mammals, lizards, and dinosaurs



Quetzalcoatlus weighed about 20 times as much as today's heaviest flying birds—an amazing 530 lb (240 kg).



Tupandactylus

TU-PAN-dac-TI-luss

The large fanlike crest of this animal was the largest of all known pterosaur crests, relative to the creature's head size. The crest was held aloft by thin bony rods.


DATE 112 mya (Early Cretaceous)

SIZE 8¼ ft (2.5 m) wingspan

FOSSIL LOCATION Brazil

HABITAT Coasts

DIET Possibly fish



Ornithocheirus

Or-NITH-oh-KEE-rus



Scientists do not know much about this creature because very few fossils have been found. After studying the fossil fragments, scientists estimate that *Ornithocheirus* had a wingspan of 33 ft (10 m) and there was a bony bump at the end of its snout, which it probably used for display.

DATE 110 mya (Early Cretaceous)

SIZE 26–33 ft (8–10 m) wingspan

FOSSIL LOCATIONS Europe and South America

HABITAT Coasts

DIET Fish

Prehistoric records

DINOSAUR RECORDS

► **Longest neck relative to body size**

In 2002, researchers working in Mongolia dug up six vertebrae, along with part of a breastbone and several leg bones of a previously unknown sauropod, which was called *Erketu ellisoni*. Based on its vertebrae, scientists estimated that its neck was 26 ft (8 m) long, making it the longest neck, relative to body size, of any known land animal.

► **Tallest dinosaur** The tallest of all dinosaurs was the sauropod *Sauroposeidon*. Although it was shorter in length than other sauropods, its neck allowed it to reach heights of up to 59 ft (18 m)—higher than any other sauropod.

► **Longest bone** The longest bone of any dinosaur belongs to a sauropod called *Ultrasaurus*. The bone was its shoulder blade and measured 8¼ ft (2.4 m) in length.

► **Brainiest dinosaur** The brainiest of all dinosaurs was *Troodon*. Relative to its body size, this theropod had the largest brain.

► **Largest skull** The ceratopsian *Pentaceratops* holds the distinction of having the largest skull of any land animal—it measured 10 ft (3 m) long.

► **Thickest skull** *Pachycephalosaurus*, an ornithischian, had the thickest skull of all dinosaurs. A bony dome on top of the skull was 8 in (20 cm) thick.

► **Longest claws** The theropod *Deinocoelurus* had the longest claws of any dinosaur. Each claw measured up to 7¼ in (19.6 cm) long.

► **Most Teeth** Of all the dinosaurs, the hadrosaur *Shantungosaurus* had the greatest number of teeth. It had more than 1,500 chewing teeth at the back of its mouth—these helped it to chew its plant food to a pulp.

PTEROSAUR RECORDS

★ **Largest pterosaur** With a wingspan of more than 36 ft (11 m), *Quetzalcoatlus* was the largest flying reptile of all time.

★ **Smallest pterosaur** *Nemicolopterus* had a wingspan of only 10 in (25 cm), making it the smallest pterosaur.

★ **Largest head crest** Relative to its body size, *Tupandactylus* had the largest head crest of all pterosaurs.

MARINE REPTILE RECORDS

- ★ **Largest ichthyosaur** *Shonisaurus* was the largest ichthyosaur. It lived in the Triassic Period and could reach lengths of up to 69 ft (21 m).
- ★ **Smallest ichthyosaur** The smallest ichthyosaur was *Chaohusaurus*. It measured only 6 ft (1.8 m) in length—the height of an average man.
- ★ **Largest plesiosaur** With a body length of more than 66 ft (20 m), *Mauisaurus* was the largest plesiosaur.
- ★ **Shortest plesiosaur** *Umoonasaurus* grew to a length of 8¼ ft (2.5 m) and was the smallest plesiosaur.
- ★ **Largest pliosaur** *Kronosaurus* was the largest of all pliosaurs. It measured up to 33 ft (10 m) in length.
- ★ **Smallest pliosaur** The smallest pliosaur skeleton found so far is that of *Leptocleidus*. This reptile was only 5 ft (1.5 m) in length.
- ★ **Largest mosasaur** *Mosasaurus* could reach lengths of up to 57¾ ft (17.6 m), making it the largest mosasaur.
- ★ **Smallest mosasaur** *Carinodens* was the smallest of all mosasaurs, although it was still quite a monster, measuring 11½ ft (3.5 m) long.

OLDEST CREATURES

- **Oldest reptile** The fossils of *Hylonomus lyelli* are older than that of any other reptile. This creature lived 312 million years ago, in the Carboniferous Period. This ancient reptile was only about 8 in (20 cm) long.
- **Oldest archosaur** *Archosaurus* lived in Russia in the Late Permian Period, around 255 million years ago, and is older than any other archosaur.
- **Oldest dinosaur** In 2011, scientists discovered fossils of *Eodromaeus*. These fossils have been dated to be more than 232 million years old—making this the oldest dinosaur found so far.
- **Oldest bird** Until recently, experts thought of *Archaeopteryx* as the oldest known bird. However, recent research by a team of Chinese scientists may change this fact. A feathered theropod called *Xiaotingia zhengi*, which lived around 155 million years ago, may have been more closely related to birds and could soon be confirmed as the earliest bird.
- **Oldest mammal** *Adelobasileus* lived in North America in the Late Triassic Period, around 220 million years ago. It is the oldest mammal.

Largest dinosaurs

Among the dinosaurs, there were some that reached gigantic proportions. Scientists are still not sure why this happened, but there are many advantages in being huge. For the largest plant-eating dinosaurs, it meant that even the most ferocious predators may have failed to bring them down. In the sauropods, gigantic digestive systems also helped to extract nutrients efficiently from plant food. Many theropods also grew larger and were able to hunt larger prey. While giant animals may live longer than small ones, they need a great amount of food, and so are vulnerable to sudden changes in climate or food supply.

LONGEST SAUROPODS

The longest sauropods were also the largest-ever land animals, therefore also the biggest dinosaurs.

- 1 **Argentinosaurus** lived in the Cretaceous Period. Scientists have found only fragments of its bones, which include vertebrae, ribs, and a thigh bone. Using these bones, they estimated that this sauropod measured between 108–134 ft (33–41 m) from head to tail.
- 2 **Supersaurus** was a Jurassic sauropod and a relative of *Apatosaurus*. Fossil remains suggest that it would have reached lengths of 108–112 ft (33–34 m).
- 3 **Sauroposeidon** is probably one of the last of the gigantic sauropods to live in North America. This Cretaceous dinosaur could reach a length of

92–112 ft (28–34 m). This sauropod is known from four neck vertebrae, which suggest that it was similar to *Brachiosaurus*.

- 4 **Futalognkosaurus** was discovered in 2000. This Cretaceous sauropod could reach 92–112 ft (28–34 m) in length, the same as *Sauroposeidon*.
- 5 **Diplodocus** lived toward the end of the Jurassic Period. It could reach lengths from 98½ ft (30 m) to 110 ft (33.5 m).
- 6 **Paralititan** is a little known Cretaceous sauropod. However, by comparing it with its relative *Saltasaurus*, scientists have estimated that it measured up to 105 ft (32 m) in length.
- 7 **Turiasaurus** was the largest sauropod in Europe, measuring more than 98½ ft (30 m) in length.

LONGEST THEROPODS

These dinosaurs were also the world's largest-ever land predators.

- 1 **Spinosaurus** was a Cretaceous theropod weighing about 7 tons and measuring 59 ft (18 m) in length.
- 2 **Carcharodontosaurus** could weigh as much as 8 tons and reach a total length of more than 46 ft (14 m).
- 3 **Giganotosaurus** lived in the late Cretaceous Period. This theropod could grow to a length of 43 ft (13 m).
- 4 **Tyrannotitan** could grow up to 40 ft (12.2 m), making it slightly larger than *Tyrannosaurus*.
- 5 **Tyrannosaurus** is the best-known of all giant theropods. It could reach 39 ft (12 m) in length and weighed more than 6 tons.
- 6 **Zhuchengtyrannus** was a cousin of *Tyrannosaurus* and lived in China during the Cretaceous Period. It could measure up to 36 ft (11 m) in length and weighed more than 6½ tons.

LONGEST ORNITHOPODS

- 1 **Shantungosaurus** lived in the Cretaceous Period. This ornithopod measured more than 52½ ft (16 m) in length.
- 2 **Lambeosaurus** is best known for its distinctive hollow crest on top of its head. It could reach a length of 49 ft (15 m).
- 3 **Edmontosaurus** was a duck-billed ornithopod, measuring up to 43 ft (13 m) long.
- 4 **Charonosaurus** was discovered in 2000. Its fossils were found near a riverbank in China. Scientists estimated that this dinosaur could grow to 43 ft (13 m) in length.
- 5 **Iguanodon** lived from the Late Jurassic Period to the Early Cretaceous Period. This ornithopod could reach lengths of more than 39 ft (12 m).
- 6 **Olorotitan** was found as a complete fossil skeleton in 2003. This dinosaur could grow to up to 39 ft (12 m).
- 7 **Saurolophus** had a body length of 39 ft (12 m).



Although some theropods were the largest predators on land, others, such as *Microraptor gui* were among the smallest of all dinosaurs.

Dinosaur discoveries

People have been discovering dinosaur bones for centuries. These bones were once thought to be of mythical creatures, and this may have given rise to the legends of dragons and giants told in different parts of the world. From the 1700s, scientists began studying these bones as evidence of animals that lived before the appearance of humans.

GREAT PALEONTOLOGISTS

Scientists who study dinosaurs and other prehistoric animals are known as paleontologists. Their research and discoveries allow us to understand prehistoric life.

► **Othniel C. Marsh (1831–99) and Edward Drinker Cope (1840–97)** were rival paleontologists who discovered many dinosaurs, including *Triceratops* and *Diplodocus*.

► **Harry Govier Seeley (1839–1909)** was a British paleontologist who classified dinosaurs into saurischians and ornithischians, based on the arrangements of their hip bones.

► **Barnum Brown (1873–1963)** was an American fossil hunter who was the first to discover fossils of *Tyrannosaurus*.

► **Elmer S. Riggs (1869–1963)** was an American paleontologist who named and described the sauropod *Brachiosaurus*, two years after its fossils were discovered.

► **Ernst Stromer von Reichenbach (1870–1952)** was a German paleontologist who named the 59-ft- (18-m-) long *Spinosaurus*.

► **Roy Chapman Andrews (1884–1960)** was an American explorer who led a number of expeditions to the Gobi Desert in Mongolia. His team discovered fossils of *Oviraptor*, *Velociraptor*, and *Proceratops*, as well as dinosaur eggs.

► **Alan Charig (1927–97)** was an American paleontologist who figured out how dinosaurs evolved from archosaurs.

► **Robert Bakker (b. 1945)** is an American paleontologist who put forward the theories that dinosaurs were warm-blooded animals and gave rise to birds.

► **John “Jack” Horner (b. 1946) and Robert Makela (1940–87)** were a team of American paleontologists who excavated dinosaur nests and found evidence that dinosaurs cared for their young.

GREAT DISCOVERIES

Since the 1600s, paleontologists have found and named more than 600 different dinosaurs. Some of the major milestones in the search for dinosaurs are given below.

🦖 **1811** Mary Anning was just 11 years old when she became the first person to find the fossils of an ichthyosaur, in the cliffs of Lyme Regis in England. She was also the first to find a plesiosaur fossil, 12 years later.

🦖 **1820** Gideon Mantell began collecting fossils of a dinosaur that he would later name *Iguanodon*. His attempts to study and reconstruct the structure and living habits of this animal spurred the scientific study of dinosaurs.

🦖 **1824** *Megalosaurus* became the first dinosaur to be named scientifically.

🦖 **1842** Sir Richard Owens coined the term “Dinosauria,” which means terrible lizard.


🦖 **1856** *Troodon* became the first American dinosaur to be given a scientific name.

🦖 **In 1861** German paleontologist Hermann von Meyer published a paper in which he described *Archaeopteryx*, which came to be considered the first bird.

🦖 **1877** Huge fossil finds in Colorado began a dinosaur rush, leading to the discovery of *Allosaurus*, *Apatosaurus*, *Diplodocus*, *Triceratops*, and *Stegosaurus*.

🦖 **1908–12** German paleontologists Werner Janensch and Edwin Hennig found fossils of *Brachiosaurus* and *Kentrosaurus* in Tanzania, Africa.

🦖 **1933–70s** Chinese paleontologist Yang Zhongjian oversaw fossil discoveries in China and named many dinosaurs, including *Lufengosaurus*, *Mamenchisaurus*, *Omeisaurus*, and *Tsintaosaurus*.



More than
486 dinosaurs
have been named
in the last
20 years.

🦖 **1979** American geologist Walter Alvarez and his father Luis Alvarez put forward the theory that the collision of an asteroid or comet with the Earth led to the extinction of the dinosaurs.

🦖 **1991** American paleontologist William Hammer discovered *Cryolophosaurus* in the Antarctic, making it the first theropod to be discovered on that continent.

🦖 **1998** Chinese paleontologists Chen Pei-ji, Dong Zhi-ming, and Zhen Shou-nan found *Sinosauropteryx*, the first dinosaur known to have a feathered body.

Glossary

3-D Having three dimensions (length, height, and depth), either in the real world, or in the virtual world of computer models or graphics.

Aetosaurus A group of plant-eating archosaurs from the Triassic Period. Aetosaurus had armored plates and spines on their backs.

Ammonites An extinct group of mollusks related to squid. They had a coiled shell and lived in Mesozoic seas.

Amphibian A vertebrate animal such as a frog that usually starts life in water as a tadpole, but turns into an air-breathing adult that lives partly on land.

Ancestor An animal or plant species from which a more recent species has evolved.

Angiosperms The plant group that contains flowering plants. This group includes broad-leaved trees and grasses.

Ankylosaurs A group of four-legged, armored, plant-eating ornithischian dinosaurs with bony plates covering the neck, shoulder, and back, and a bony club at the end of the tail.

Archosaurs A group of prehistoric reptiles that include the dinosaurs, pterosaurs, and crocodylians and their relatives. The archosaurs had a distinctive cavity in the skull on each side of the snout between the eye and the nostril. They evolved around 255 million years ago.

Arid Describes a dry climate or place.

Asteroid A large rocky object that orbits the Sun—bigger than a meteoroid but smaller than a planet.

Asymmetrical Having unequal parts.

Binocular vision Viewing a scene or object with two eyes. This enables an animal to judge distance to the scene or object.

Bipedal Describes animals that walk on two hind limbs.

Bird A feathered theropod dinosaur capable of powered flight or an animal descended from such a flying ancestor.

Brackish Water that is saltier than fresh water, but less salty than ocean water.

Cambrian The first period of the Paleozoic Era, lasting from 542 to 488 million years ago. Most of the main animal groups evolved in this period.

Camouflage Colors or patterns that help an animal to blend in with its surroundings.

Carnivore An animal that eats only meat.

Cenozoic The era of time that followed the Mesozoic. It began 65 million years ago and extends up to the present day. Cenozoic literally means “new animal life.”

Ceratopsians A group of four-legged, horned, plant-eating ornithischian dinosaurs, such as *Triceratops*, with a bony frill at the back of the skull.

CGI Short for computer-generated imagery, CGI is any illustration created using a computer, including 3-D models and animation sequences.

Cold-blooded Describes an animal whose body temperature rises and falls along with the temperature of its surroundings.

Colony A group of separate individuals working together, as in an ant or bird colony, or animals joined by strands of living tissues as in the case of corals.

Coniferous Cone-bearing trees, such as pine or fir.

Coprolite Fossilized dung of an animal.

Cretaceous Period The third period of the Mesozoic Era, lasting from 145 to 65 million years ago. An asteroid or comet hit Earth at the end of this period, triggering the demise of the dinosaurs.

Crocodylians The group that includes living crocodiles and alligators and their immediate ancestors. They and their extinct relatives are known as the crocodylomorphs and belong to the archosaur group of reptiles.

Crurotarsans A group of archosaurs that included the crocodylomorphs and other related reptiles, such as the rauisuchians and aetosaurus.

Cycad A tropical or subtropical plant that bears its seeds in large cones, but has a crown of leaves like a tree fern or palm.

Cynodonts A group of mammal-like reptiles with mammal-like teeth, which include canines, incisors, and molars. They included the ancestors of mammals.

Dicynodonts A group of plant-eating mammal-like reptiles with two tusks and a blunt beak.

Dinosaurs A group of archosaurs that evolved in the Triassic Period and dominated life on land in the Jurassic until they went extinct at the end of the Cretaceous. The birds are their direct descendants.

Ecosystem A collection of organisms living together in the same habitat, together with their environment.

Embryo An animal or plant in an early stage of development from an egg or a seed.

Environment The natural surroundings in which an animal or plant lives.

Era A unit of time that is further divided into periods.

Evolution The gradual change in living organisms that occur over many generations. Evolution

may result in new species. Dinosaurs evolved from archosaur ancestors and birds evolved from feathered theropods.

Extinction The dying out of a plant or animal species. Extinction can happen naturally as a result of competition between species, changes in the environment, natural disasters (such as an asteroid or comet striking the Earth), or other factors.

Ferns A group of nonflowering plants that reproduce by spores rather than by seeds.

Fossil The remains of a dead organism, preserved in rock. Teeth and bones are more likely to form fossils than softer body parts, such as muscles and internal organs.

Fossilization The process by which dead organisms turn into fossils.

Genus (plural, Genera) In the classification of living organisms, a group of closely related species. The theropod *Tyrannosaurus rex* belongs to the genus *Tyrannosaurus*.

Ginkgo One of a group of nonflowering plants that grows into a tall tree with triangular leaves.

Gondwana One of the two continents that Pangaea split into at the end of the Triassic Period.

Gymnosperms One of the two main types of land plant that produce seeds. This group includes cycads, ginkgos, and conifers, such as pine and fir.

Hadrosaurs A group of large, bipedal and quadrupedal ornithopods from the Late Cretaceous Period known as duck-billed dinosaurs. They had a ducklike bill and fed on plants.

Herbivore An animal that eats only plants.

Hominids A group of primate mammals that includes humans, chimpanzees, and gorillas, together with their extinct close relatives. It does not include orangutans, gibbons, and monkeys.

Hypsilophodonts A group of fast-moving, bipedal ornithopods.

Ichthyosaurs A group of dolphinlike, predatory marine reptiles that were common in the Mesozoic Era. They had large eyes, pointed heads, and sharklike fins and tails.

Iguanodonts A group of ornithopod dinosaurs that ranged from small to large. They all had horselike faces. The group was named after an ornithopod called *Iguanodon*.

Invertebrate An animal without a backbone.

Jurassic Period The second period of the Mesozoic Era, lasting from 200 to 145 million years ago. During the Jurassic Period, dinosaurs dominated the land, the first birds evolved, and mammals began to spread out.

Keelbone The enlarged, deepened breastbone of a bird that anchors the large flight muscles. All modern birds have one, but some early birds did not.

Keratin The substance that forms skin, hair, horns, and nails or hooves of an animal.

Laurasia One of the two continents that were formed after Pangaea split at the end of the Triassic.

Lava The molten rock expelled by an erupting volcano or the solid rock that forms after cooling.

Mammals A group of warm-blooded vertebrates that feed their young on milk and whose skin is covered in hair or fur. Mammals evolved from cynodont ancestors in the Triassic Period.

Mesozoic Era The era lasting from 252 to 65 million years ago, containing the Triassic, Jurassic, and Cretaceous periods.

Metriorhynchids A family of fierce, marine crocodylian predators that had a streamlined body.

Mollusks A large group of invertebrate species including slugs, snails, clams, octopuses, and squid. Many mollusks produce hard shells that fossilize easily, making mollusk fossils common.

Mosasaurs A group of large aquatic lizards with paddle-shaped limbs and a tail flattened from side to side. They hunted fish and other marine creatures in the Cretaceous Period.

Nesting colony A colony of animals, such as birds and dinosaurs, that is formed when the animals gather in the same area to build nests and care for their eggs and young.

Nodosaurs A group of four-legged, armored, plant-eating dinosaurs that had bony plates on their backs and spines on the sides. They were related to the ankylosaurs.

Nothosaurs A group of predatory reptiles that lived in the Triassic seas. They had four legs with webbed feet and may have bred on shores like seals.

Omnivore An animal that eats both plants and other animals.

Ornithischians One of the two main dinosaur groups. In ornithischians, the pelvis (hip bone) was arranged as in a bird. Ornithischians included stegosaurs, ankylosaurs, ceratopsians, ornithopods, and pachycephalosaurs.

Ornithomimids A group of theropods that resembled ostriches. They were known as ostrich dinosaurs.

Ornithopods A group of plant-eating, mainly bipedal, ornithischians with long hind limbs. This group includes the hypsilophodonts, iguanodonts, and hadrosaurs.

Oviraptorosaurs A group of theropods with parrotlike skulls and feathered bodies.

Pachycephalosaurs A group of bipedal dinosaurs with thick, domed skulls.

Paleontologist A scientist who studies the fossil remains of plants and animals.

Paleontology The scientific study of fossilized plants and animals.

Paleozoic The era of time that came before the Mesozoic Era. It lasted from 542 to 252 million years ago. Paleozoic means “ancient animal life.”

Pangaea The supercontinent that existed in the Paleozoic and Mesozoic Eras.

Period A unit of time that is a division of an era—the Triassic Period is part of the Mesozoic Era.

Pigments Chemicals that give color to an organism. Giving color may be the main function of the pigment or the coloration may be incidental.

Placenta The organ that forms inside a pregnant mammal partly from the embryo and partly from the wall of the mother’s womb, or uterus. The placenta allows exchange of nutrients and waste

between the mother and developing young of placental mammals. *Eomaia* was one of the first placental mammals.

Plesiosaurs A group of meat-eating sea reptiles from the Jurassic and Cretaceous periods that swam in the seas using flipperlike limbs. Many had a snakelike neck and a small head.

Pliosaurus A group of plesiosaurs that had a short muscular neck and a large head with crocodilelike teeth. They were among the most formidable of all marine predators.

Precambrian The time between the formation of the Earth and the beginning of the Cambrian Period.

Predator An animal that hunts, kills, and eats other animals for food.

Primates A group of mammals that includes lemurs, monkeys, apes, and humans.

Primitive At an early stage of evolution.

Prosauropods A group of early, mainly plant-eating saurischian dinosaurs. They were the ancestors of the giant, long-necked sauropods.

Psittacosaur A group of bipedal ceratopsians from the Cretaceous Period. Psittacosaur had deep, parrotlike beaks that they used to eat plants.

Pterosaur A group of archosaur reptiles that were capable of powered flight. They had batlike wings made of skin. Some of these were the largest creatures ever to fly.

Pubis One of the three bones that forms the pelvis of an animal.

Quadrupedal Describes an animal that walks on four limbs.

Rauisuchians A group of archosaur reptiles with upright limbs that lived in the Triassic. Many rauisuchians were dinosaurlike.

Reptile A group of cold-blooded vertebrates with scaly skin that typically live on land and reproduce by laying eggs. Lizards, snakes, turtles, and crocodiles are reptiles.

Saurischians One of the two main dinosaur groups. In saurischians, the pelvis was arranged as in a lizard. Saurischians include the predatory theropods and the sauropodomorphs.

Sauropodomorphs A group of plant-eating saurischians, including the prosauropods and sauropods.

Sauropods A group of gigantic, long-necked saurischians that included some of the largest animals ever to walk on the Earth.

Scutes Bony plates with a covering made of horn, set in the skin of certain reptiles. Scutes help form body armor in these reptiles.

Sediment Material such as sand and mud deposited by wind, water, or ice.

Serrated Saw-toothed, like a steak knife.

Skull The head's bony framework that protects the brain, eyes, ears, and nasal passages.

Stegosaurs A group of four-legged, plant-eating ornithischian dinosaurs with rows of tall bony plates and spines running down their backs and tails.

Stromatolites Large, hard, domelike structures in shallow seawaters, made of thin layers of particles built up by the action of generations of microorganisms, such as cyanobacteria.

Stromatolites were very common in Precambrian times.

Temperate Describes a climate that is neither very hot nor very cold.

Territory The part of an animal's habitat that it defends from rival animals, usually of its own species.

Tetrapods Vertebrates with four limbs (arms, legs, or wings). All amphibians, reptiles, mammals, and birds are tetrapods. All tetrapods evolved from a fishlike ancestor.

Theropods A group of meat-eating dinosaurs. All theropods were predatory. They typically had sharp teeth and claws and ranged in size from the tiny *Microaptor* to the colossal *Tyrannosaurus*.

Trace fossils The signs of prehistoric creatures or their activities rather than remains of the creatures themselves, preserved in rock. Trace fossils include footprints, bite marks, droppings, and eggs.

Triassic Period The first period of the Mesozoic Era, lasting from 252 to 200 million years ago. Dinosaurs evolved in the Triassic Period.

Tropical Describes the region that spans the equator. It is a hot region with plenty of rainfall, resulting in the growth of tropical rainforests.

Tyrannosaurids A group of large theropods with short arms and two-fingered hands. They were named after *Tyrannosaurus*.

Vertebrae The bones that make up the backbone, or spine, of an animal.

Vertebrates Animals with a spinal column, or backbone.

Warm-blooded Describes an animal that maintains a constant internal body temperature. Mammals and birds are warm-blooded and at least some dinosaurs were, too. Their body temperature does not change with the temperature of their surroundings.

Wingspan The distance from the tip of one wing of an animal to the tip of the other when both wings are outstretched.

Index

A

Acanthostega 14
Adelobasileus 143
 aetosaurs 106–7
Albertosaurus 46
Allosaurus 42, 147
 Alvarez, Walter and Luis 147
Alxasaurus 17
Amargasaurus 62
 ammonites 23
Amphibamus 5
Anchisaurus 57
 Andrews, Roy Chapman 146
 ankylosaurs 7, 16–17, 80–3
Ankylosaurus 16–17, 80
 Anning, Mary 147
Anomalocaris 4
Anurognathus 138
Apatosaurus 28, 61, 147
Archaeopterus 5
Archaeopteryx 6, 19, 54, 143, 147
 archosaurs 15, 103, 106–13, 136, 143, 146
Archosaurus 143
Argentinosaurus 67, 144
 armored dinosaurs 16, 72–83
 asteroids 7, 20–1, 147

B

Bakker, Robert 146
Barosaurus 16, 62–3, 64–5
Barsboldia 44–5
Baryonyx 25, 36, 40

birds
 early 54–5, 146
 evolution 13, 18–19
 oldest 19, 143, 147
 bones
 fossil 24–5, 146–7
 largest 142
 bony growths 72, 78, 80, 96, 97, 98
Brachiosaurus 11, 68–9, 146, 147
Brachylophosaurus 95
 brain size 142
Brontosaurus 61
 Brown, Barnum 146

C

Camarasaurus 70
 Cambrian Period 4
Camptosaurus 90
 Carboniferous Period 5
Carcharodontosaurus 43, 145
Carinodens 143
Caudipteryx 49, 53
 Cenozoic Era 7
Centrosaurus 98
 ceratopsians 7, 13, 17, 98–101
Ceratosaurus 37
 CGI (computer-generated imagery) 31
Chaohusaurus 143
 Charig, Alan 146
Charonosaurus 145
Chasmosaurus 100–1
 Chen Pei-ji 147

Chicxulub crater 21
Citipati 49, 52–3
 claws
 fossil 25
 longest 142
 thumbs 56
 climate change 20
 clubs, bony 80, 83
Coelophysis 9, 36
 coloration 31, 33
 comets 7, 20–1, 147
Compsognathus 46
 Cope, Edward Drinker 146
 coprolite 28
Corythosaurus 31, 92–3
 crests 92–5, 142
 Cretaceous Period 6, 7, 12–13
 crocodilians 15, 103, 106–7, 110, 111
 crurotarsans 106–7
Cryolophosaurus 38–9, 147
Cryptoclidus 126
 cynodonts 103, 114–15
Cynognathus 115

D

Dakosaurus 110
Deinonychus 36, 51
Deinosuchus 107, 111
Desmatosuchus 107
 Devonian Period 5
Dicraeosaurus 63
 dicynodonts 114–15
Dilophosaurus 38
Dimetrodon 5
Dimorphodon 137

dinosaurs 32–101
 discoveries 146–7
 evolution 6, 9, 14–15, 106
 extinction 7, 20–1, 147
 fossils 22–9
 largest 144–5
 neighbors 102–17
 reconstruction 30–1
 records 142–3
 types 16–17
Diplodocus 60, 144, 146,
 147
 Dong Zhi-ming 147
 dryosaurids 85
Dryosaurus 85
 duck-billed dinosaurs 12, 13,
 84, 92–5

E

Edmontonia 78
Edmontosaurus 26, 94–5,
 145
Effigia 103, 107, 108–9,
 112–13
Efraasia 56
 eggs
 evolution 14
 fossilized 28–9, 53, 146
Einosaurus 100–1
Elasmosaurus 127
 embryo fossils 29
Eocursor 35
Eodromaeus 143
Eomaia 103, 117
Eoraptor 6, 35
Eosimias 7
 eras 4
Erketu ellisoni 142
Eudimorphodon 9, 137

Euoplocephalus 81, 82–3
 extinction 7, 8, 20–1, 144,
 147

F

feathers
 evolution 19
 fossilized 27, 147
 ferns 8
 fish 14
 flowering plants 12
 footprints 29, 60–1
 fossils
 bones 24–5
 discoveries 146–7
 exceptional 26–7
 formation of 22–3
 trace 23, 28–9
 frills 98
Futalognkosaurus 144

G

Gallimimus 25, 48
Gargoyleosaurus 79
Gastonia 78
Giganotosaurus 43, 145
Gojirasaurus 35
 Gondwana 10, 12
Gryposaurus 25
Guanlong 47

H

Hadrosaurus 92
 hadrosaurs 44, 84, 92–5
 Hammer, William 147
 Hennig, Edwin 147
Herrerasaurus 34

Heterodontosaurus 84
 hominids 6, 7
 Horner, John “Jack” 146
 horns 16, 98
Huayangosaurus 76–7
Hylonomus lyelli 143
Hyperodapedon 9, 104–5
Hypsilophodon 86
 hypsilophodonts 84, 86, 90

I

Iberomesornis 19
Ichthyornis 55
 ichthyosaurs 10, 28, 124–5,
 143, 147
Ichthyosaurus 10, 125
Iguanodon 24, 29, 91, 145,
 147
 iguanodonts 84, 90–1
 invertebrates 4

JK

Janensch, Werner 147
 jaws 36
 Jurassic Period 6, 10–11, 33
Kayentachelys 120
Kentrosaurus 76–7, 147
 keratin 75
Kronosaurus 129, 143

L

Lagosuchus 108
Lambeosaurus 93, 145
Lariosaurus 122
 Laurasia 10, 12
Leaellynasaura 86
Leptocleidus 143

Lesothosaurus 85, 88–9
Liaoxiornis 13
Liliensternus 37
Liopleurodon 128
Lufengosaurus 58–9, 147
Lystrosaurus 115

M

Maiasaura 13, 92
 Makela, Robert 146
Mamenchisaurus 6, 68, 147
 mammals 7, 8, 13, 21, 103
 early 116–17
 Mantell, Gideon 24, 147
 marine reptiles *see* sea
 reptiles
 Marsh, Othniel C. 146
 mass extinctions *see*
 extinction
Massopondylus 59
Maisaurus 143
 meat-eating dinosaurs 11, 16,
 34, 36
Megalosaurus 91, 147
Megazostrodon 117
 Mesozoic Era 6, 7, 10, 16,
 103, 135
 metriorhynchids 110
 Meyer, Hermann von 147
Microraptor 19, 50
Minmi 81
Mixosaurus 125
 monkey puzzle trees 10
Monolophosaurus 38
Morganucodon 116
 mosasaurs 13, 119, 132–3, 143
Mosasaurus 13, 132, 143
 movement 31
Muttaborrasaurus 16, 87

N

necks 142
Nemegtbaatar 116
Nemegtosaurus 71
Nemicolopterus 142
 Neogene Period 7
 nests 29, 53, 146
 nodosaurs 78–9
 nothosaurs 119, 122–3
Nothosaurus 123

O

Odontochelys 121
Olorotitan 145
Ornaisaurus 147
 omnivores 34, 36
 ornithischians 17, 146
Ornithocheirus 141
Ornitholestes 48
Ornithomimus 48–9
 ornithopods 16, 17, 84–95,
 145
Ornithosuchus 109
Oviraptor 29, 53, 146
 oviraptorosaurs 53
 Owens, Sir Richard 147

P

pachycephalosaurs 17,
 96–7
Pachycephalosaurius 97,
 142
Pachypleurosaurus 122
 Paleogene Period 7, 21
 paleontologists 24, 146–7
 Paleozoic Era 4–5
Panderichthys 14
 Pangaea 8, 10
Paralititan 144
Parasaurolophus 95
Parasuchus 15, 108
Pentaceratops 100, 142
 periods 4
 Permian Period 5
Peteinosaurus 138
Phenacodus 21
 phytosaurs 108
Placerias 114
 placodonts 120
Placodus 120
 plant-eating dinosaurs 16, 34,
 36, 56
 plants 5, 8, 10, 12, 33
Pleuromeia 58
 plates 72, 78, 80
 plesiosaurs 10, 119, 126–7,
 143, 147
Plesiosaurus 127
Pleuromeia 8–9
Plioplatecarpus 133
 pliosaurs 128–31, 143
Polacanthus 26
Postosuchus 15, 102–3, 107,
 108–9
 Precambrian Era 4
 primates 7
Proceratops 146
Proceratosaurus 47
 prosauropods 17, 56–9
Protoceratops 99
Psittacosaurus 98
 pterodactyls 137
Pterodactylus 11, 135,
 138–9
 pterosaurs 9, 11, 106,
 134–41, 142
 Purgatoire River track site 61

QR

Quaternary Period 7
Quetzalcoatlus 134–5, 140, 142
 raurisuchians 108–9, 112–13
 reconstruction 30–1
 records, prehistoric 142–3
 Reichenbach, Ernest Stromer von 146
 reptiles 5, 14, 15, 143
Rhabdodon 90
Rhamphorhynchus 136–7
Rhomaleosaurus 129, 130–1
 rhynchosaurs 9, 103, 104–5
Rhynchosaurus 104–5
 Riggs, Elmer S. 146

S

Saltasaurus 66–7
 saurischians 17, 146
Saurolophus 145
Sauropelta 79
 sauropodomorphs 17
 sauropods 6, 11, 16, 17, 60–71, 144
Sauroposeidon 142, 144
Scelidosaurus 74–5
Scutellosaurus 72
 sea reptiles 10, 118–32, 143
 Seeley, Harry Govier 146
Shangtungosaurus 142, 145
Shonisaurus 124, 143
Shunosaurus 70
Simosuchus 111
Sinoconodon 117
Sinornithosaurus 18, 27
Sinosauropteryx 18, 147
Sinraptor 43

skeletons 24, 25, 30
 skin, fossilized 23, 26
 skulls
 fossil 25
 largest/thickest 142
Sphenosuchus 88–9, 110
 spines and spikes 16, 72, 78, 97
Spinosaurus 41, 145, 146
Stagonolepis 106
Stegoceras 96
 stegosaurs 17, 72–7
Stegosaurus 73, 147
 stromatolites 4
Styracosaurus 17, 101
Sucomimus 40
Supersaurus 144

T

tails 80, 83, 136–7
Tarbosaurus 44–5
 teeth 16, 56, 114, 142
Temnodontosaurus 125
Tempyska 12
Tenontosaurus 86–7
Terrestriusuchus 110
 tetrapods 5, 14
Thecodontosaurus 57
 theropods 6, 11, 17, 18, 19, 36–53, 145
Thrinaxodon 115
Titanosaurus 66
 trace fossils 23, 28–9
 Triassic Period 6, 8–9, 34, 116
Triceratops 13, 98, 99, 146, 147
Troodon 50–1, 142, 147
Tsintaosaurus 147
Tuojiangosaurus 77

Tupandactylus 141, 142
Turiasaurus 144
 turtles 119, 120–1
 tusks 114
Tyrannosaurus 36, 46–7, 145, 146
Tyrannotitan 145

UVW

Ultrasaurus 142
Umoonasaurus 143
Vegavis 55
Velociraptor 51, 146
 vertebrae 142
 vertebrates 4, 5
 volcanic activity 20, 21
Vulcanodon 63
Westlothiana 14
Williamsonia 10

XYZ

Xiaotingia 54, 143
Xiphactinus 23
 Yang Zhongjian 147
Zalambdalestes 13
 Zhen Shou-nan 147
Zhuchengtyrannus 145

Acknowledgments

Dorling Kindersley would like to thank: Stewart Wild for proofreading; Helen Peters for indexing; David Roberts and Rob Campbell for database creation; Claire Bowers, Fabian Harry, Romaine Werblow, and Rose Horridge for DK Picture Library Assistance; Tanveer Abbas Zaidi for CTS assistance; and Tanya Mehrotra and Mahipal Singh for design assistance.

The publishers would also like to thank the following for their kind permission to reproduce their photographs:

(Key: a-above; b-below/bottom; c-centre; f-far; l-left; r-right; t-top)

1 Dorling Kindersley: Peter Minister, Digital Sculptor (c). **2-3 Dorling Kindersley:** Jon Hughes (c). **4 Nature.com:** Doug Perrine (b). **6-7 Science Photo Library:** MARK GARLICK (bc). **6 Dorling Kindersley:** Jon Hughes (r). **7 Getty Images:** Dan Kitwood (tr). **8 Dorling Kindersley:** Andy Crawford / David Donkin - modelmaker (tl). **9 Dorling Kindersley:** Jon Hughes (ca); Peter Minister, Digital Sculptor (br). **10 Dorling Kindersley:** Andy Crawford / David Donkin - modelmaker (tl). **11 Dorling Kindersley:** Andrew Kerr (b); Peter Minister, Digital Sculptor (tl). **Getty Images:** Siri Stafford / Riser (b/background). **12 Dorling Kindersley:** Andy Crawford / David Donkin - modelmaker (tl). **13 Corbis:** Inspirostock (br/background). **Dorling Kindersley:** Peter Minister, Digital Sculptor (br). **14 Dorling Kindersley:** Natural History Museum, London (b); The Oxford University Museum of Natural History (cr). **16 Dorling Kindersley:** Peter Minister, Digital Sculptor (bl). **17 Dorling Kindersley:** Tim Ridley / Robert L. Braun - modelmaker (bc). **18 Corbis:** Grant Delin / Outline Gallery (bc). **Reuters:** Mike Segal (r). **19 Dorling Kindersley:** Peter Minister, Digital Sculptor (cr). **Science Photo Library:** Christian Darkin (tl). **20 Science Photo Library:** MARK GARLICK (bc). **21 Alamy Images:** Yoghesh More / ephotocorp (cl). **Dorling Kindersley:** Bedrock Studios / Jon Hughes (br). **Science Photo Library:** D. Van Ravenswaay (tl). **23 Dorling Kindersley:** James Stevenson / Donks Models - modelmaker (br). **24 Dorling Kindersley:** Natural History Museum, London. **25 Dorling Kindersley:** Royal Tyrrell Museum of Palaeontology, Alberta, Canada (tr). **26 Dorling Kindersley:** Natural History Museum, London (bl). **The Natural History Museum, London:** (tr). **27 Reuters:** Mike Segar (r). **29 Corbis:** Louie Psihoyos / Terra (tr). **Dorling Kindersley:** Natural History Museum, London (bl). **30 Corbis:** Louie Psihoyos / Terra. **32 Corbis:** Philippe Widling / Design Pics (background). **Dorling Kindersley:** Peter Minister, Digital Sculptor. **33 Corbis:** Kevin Schafer (br/background). **Dorling Kindersley:** Peter Minister, Digital Sculptor (br). **36 Corbis:** Randall Levensaler Photography / Aurora Photos (br/background). **Dorling Kindersley:** Natural History Museum, London (tl). **Peter Minister, Digital Sculptor (br).** **38-39 Corbis:** Philippe Widling / Design Pics (background). **Dorling**

Kindersley: Peter Minister, Digital Sculptor. **38 Dorling Kindersley:** Andy Crawford / Robert L. Braun - modelmaker (tl). **40 Dorling Kindersley:** Jon Hughes (tl); Gary Ombler / Jonathan Hatley - modelmaker (b). **42 Corbis:** Gary Weathers / Tetra Images (background). **Dorling Kindersley:** Peter Minister, Digital Sculptor. **43 Dorling Kindersley:** Gary Ombler (bl). **46 Corbis:** Mitsushi Okada / amanaimages (bl/background). **46-47 Corbis:** Randall Levensaler Photography / Aurora Photos (bc/background). **Dorling Kindersley:** Peter Minister, Digital Sculptor (c). **47 Dorling Kindersley:** Natural History Museum, London (br). **48 Dorling Kindersley:** Royal Tyrrell Museum of Palaeontology, Alberta, Canada (b). **48-49 Dorling Kindersley:** Royal Tyrrell Museum of Palaeontology, Alberta, Canada (tl). **49 Corbis:** Owen Franken (r/background). **Dorling Kindersley:** Peter Minister, Digital Sculptor (r). **50 Science Photo Library:** Christian Darkin (b). **51 Dorling Kindersley:** Jon Hughes (bl); Peter Minister, Digital Sculptor (br). **52-53 Corbis:** Owen Franken (background). **Dorling Kindersley:** Peter Minister, Digital Sculptor. **54 Corbis:** moodboard (b/background). **Dorling Kindersley:** Peter Minister, Digital Sculptor (br). **55 Dorling Kindersley:** Jon Hughes (bl); Natural History Museum, London (tc); Peter Minister, Digital Sculptor (tl). **56 Dorling Kindersley:** Jon Hughes (br); Institute of Geology and Palaeontology, Tubingen, Germany (cl, bl). **61 Alamy Images:** Rob Walls (tl). **Corbis:** Louie Psihoyos / Science Faction (tr). **62-63 Dorling Kindersley:** Peter Minister, Digital Sculptor (tl). **Getty Images:** Siri Stafford / Riser (t/background). **62 Getty Images:** Jeffrey L. Osborn / National Geographic (br). **64-65 Dorling Kindersley:** Peter Minister, Digital Sculptor. **Getty Images:** Siri Stafford / Riser (background). **66-67 Corbis:** Alan Traeger (b/background). **Dorling Kindersley:** Peter Minister, Digital Sculptor (bl). **68 Dorling Kindersley:** Jon Hughes (tl). **68-69 Dorling Kindersley:** Andrew Kerr. **Getty Images:** Siri Stafford / Riser (r/background). **68 Dorling Kindersley:** Jon Hughes (tl). **70 The Natural History Museum, London:** (br). **73 Dorling Kindersley:** Peter Minister, Digital Sculptor. **Getty Images:** Holger Spiering (background). **74-75 Dorling Kindersley:** Peter Minister, Digital Sculptor. **Getty Images:** Oliver Strewé / Photographer's Choice (background). **77 Dorling Kindersley:** Leicester Museum (r). **80 Dorling Kindersley:** Jon Hughes (b). **82-83 Corbis:** John Carnemolla (background). **Dorling Kindersley:** Peter Minister, Digital Sculptor. **84 Corbis:** Kevin Schafer (br/background). **Dorling Kindersley:** Peter Minister, Digital Sculptor (br). **85 Corbis:** Paul A. Souders (t/background). **Dorling Kindersley:** Peter Minister, Digital Sculptor (tl). **88-89 Corbis:** Paul A. Souders (background). **Dorling Kindersley:** Peter Minister, Digital Sculptor. **91 Dorling Kindersley:** Jon Hughes. **94-95 Dorling Kindersley:** Peter Minister, Digital Sculptor. **Getty Images:** Panoramic Images (background).

95 Dorling Kindersley: Peter Minister, Digital Sculptor (br); Royal Tyrrell Museum of Palaeontology, Alberta, Canada (tr). **96 Dorling Kindersley:** Royal Tyrrell Museum of Palaeontology, Alberta, Canada (br). **97 Corbis:** Raelius Images (background). **Dorling Kindersley:** Peter Minister, Digital Sculptor. **98 Dorling Kindersley:** Natural History Museum, London (bl, cl, br, tl). **99 Corbis:** Inspirostock (br/background). **100-101 Getty Images:** De Agostini Picture Library (tc). **104-105 Dorling Kindersley:** Institute of Geology and Palaeontology, Tubingen, Germany (b). **105 Dorling Kindersley:** Natural History Museum, London (tl). **107 Dorling Kindersley:** Jon Hughes (br). **108 Dorling Kindersley:** Jon Hughes (tl). **114 Dorling Kindersley:** Natural History Museum, London (cl). **115 Dorling Kindersley:** Natural History Museum, London (bl). **122 Dorling Kindersley:** Royal Tyrrell Museum of Palaeontology, Alberta, Canada (bl). **123 Dorling Kindersley:** David Peart (background). **125 Dorling Kindersley:** Jon Hughes (br). **126 Dorling Kindersley:** Hunterian Museum (University of Glasgow) (b). **129 Corbis:** Mark A. Johnson (tr/background). **Dorling Kindersley:** Peter Minister, Digital Sculptor (tr). **130-131 Corbis:** Mark A. Johnson (background). **Dorling Kindersley:** Peter Minister, Digital Sculptor. **135 Dorling Kindersley:** Peter Minister, Digital Sculptor (bc). **136 Dorling Kindersley:** Jon Hughes (tl). **137 Dorling Kindersley:** Jon Hughes (bl); Peter Minister, Digital Sculptor (tr); Robert L. Braun - modelmaker (br). **Getty Images:** DAJ (tr/background). **138-139 Dorling Kindersley:** Peter Minister, Digital Sculptor. **Getty Images:** DAJ (background). **141 Dorling Kindersley:** Peter Minister, Digital Sculptor (tr).

Jacket images: *Front:* Dorling Kindersley: The American Museum of Natural History (bl), Graham High at Centaur Studios - modelmaker tl, cr/ (*Brachiosaurus*), cb, Jon Hughes tr, cl/ (*Argentinosaurus*), bl/ (*Velociraptor*), Jon Hughes / Bedrock Studios tr, tl/ (*Scutosaurus*), cla, Graham High - modelmaker tl/ (*Pentaceratops*), Jonathan Hatley - modelmaker cr/ (*Baryonyx*), Robert L. Braun - modelmaker ca, cr/ (*Dilophosaurus*), cr/ (*Camotaurus*), br, Centaur Studios - modelmakers c/ (*Triceratops*), Leicester Museum (bl/ (*Tuojiangosaurus*), Natural History Museum, London cr/ (*Archaeopteryx*), bl, bl/ (*Psittacosaurus*), br/ (*Triceratops*), Peter Minister, Digital Sculptor (bl/ (*Deinonychus*), c, Tim Ridley / Robert L. Braun - modelmaker br/ (*Styracosaurus*), Royal Tyrrell Museum of Palaeontology, Alberta, Canada tr/ (skeleton), cr, cr; *Back:* Dorling Kindersley: Jon Hughes (cl), Jonathan Hatley - modelmaker cl, Royal Tyrrell Museum of Palaeontology, Alberta, Canada cl; *Spine:* Dorling Kindersley: Peter Minister, Digital Sculptor t.

All other images © Dorling Kindersley

For further information see: www.dkimages.com